


Fisheries Management Report No. 4
of the

ATLANTIC STATES MARINE
FISHERIES COMMISSION



FISHERY
MANAGEMENT
PLAN
FOR
SPOTTED
SEATROUT

October 1984

QL638
.S34M383
1984

FISHERY MANAGEMENT PLAN
FOR THE
SPOTTED SEATROUT (Cynoscion nebulosus) FISHERY

Prepared by
Linda P. Mercer, Project Leader

North Carolina Department of Natural Resources
and Community Development
Division of Marine Fisheries
Morehead City, NC 28557

Special Scientific Report No. 43

October 1984

This plan was prepared in cooperation with the Sciaenid Technical Committee and Sciaenid Board, as part of the Interstate Fishery Management Program administered by the Atlantic States Marine Fisheries Commission.

Funds provided by Southeast Region, National Marine Fisheries Service, National Oceanic and Atmospheric Administration under Cooperative Agreement No. SF-13 and by Northeast Region, National Marine Fisheries Service, National Oceanic and Atmospheric Administration under Cooperative Agreement No. NA-80-FA-H-00017.

200 copies of this public document were printed at
a cost of \$710.39 or \$3.55 per copy.

638-534 M383 1984

1.0 EXECUTIVE SUMMARY

The purpose of this fishery management plan is to manage the spotted seatrout fishery in the territorial sea of the Atlantic Ocean from Maryland to Florida. The goal of the plan is to perpetuate the spotted seatrout resource in fishable abundance throughout its range and generate the greatest possible economic and social benefits from its harvest and utilization over time.

Plan objectives and management measures are directed toward alleviating the following problems:

1. Lack of biological, social and economic data needed to define optimum yield.
2. Protection of adequate spawning stock in years of severe environmental conditions.
3. Recreational-commercial conflicts.

Management objectives designed to address the above problems are:

1. Attain over time optimum yield.
2. Maintain a spawning stock sufficient to minimize the possibility of recruitment failure.
3. Promote conservation of the stocks in order to reduce inter-annual variation in availability and increase yield per recruit.
4. Promote the collection of economic, social, and biological data required to effectively monitor and assess management efforts relative to the overall goal.
5. Promote research that improves understanding of the biology and fisheries of spotted seatrout.
6. Promote harmonious use of the resource among various components of the fishery through the coordination of management efforts among the various political entities having jurisdiction over the spotted seatrout resource.
7. Promote determination and adoption of standards of environmental quality and provide habitat protection necessary for the maximum natural production of spotted seatrout.

Management measures include a minimum size limit of 12 inches total length with comparable mesh size regulations in directed fisheries, and data collection for stock assessment and to monitor the status of the fisheries. High research priorities include stock identification, mortality estimates and yield modeling, habitat requirements, effects of environmental factors on stock size, development of a prerecruit index, mesh size selectivity, and social and economic analyses.

2.0 CONTENTS

	Page
1.0 EXECUTIVE SUMMARY	i
2.0 CONTENTS	iii
3.0 INTRODUCTION.	1
3.1 Development of the Plan.	1
3.2 Problems Addressed by the Plan	1
4.0 DESCRIPTION OF STOCKS	2
4.1 Species Distribution	2
4.2 Abundance and Present Condition	2
4.3 Ecological Relationships	2
4.4 Estimate of Maximum Sustainable Yield	3
4.5 Probable Future Condition	3
5.0 DESCRIPTION OF HABITAT	3
5.1 Condition of the Habitat	3
5.2 Habitat Areas of Particular Concern	4
5.3 Habitat Protection Programs.	4
6.0 FISHERY MANAGEMENT JURISDICTION, LAWS, AND POLICIES . . .	6
6.1 Management Institutions	6
6.2 Treaties and International Agreements	6
6.3 Federal Laws, Regulations, and Policies	7
6.4 State Laws, Regulations, and Policies	7
6.5 Local and Other Applicable Laws, Regulations, and Policies	7
7.0 DESCRIPTION OF FISHING ACTIVITIES	7
7.1 History of Exploitation	7
7.2 Domestic Commercial and Recreational Fishing Activities	8
7.3 Foreign Fishing Activities	8

	Page
8.0 DESCRIPTION OF ECONOMIC CHARACTERISTICS OF THE FISHERY	8
8.1 Domestic Harvesting Sector	8
8.2 Domestic Processing Sector	12
8.3 International Trade	12
9.0 DESCRIPTION OF BUSINESSES, MARKETS, AND ORGANIZATIONS ASSOCIATED WITH THE FISHERY	12
9.1 Relationship Among Harvesting and Processing Sectors	12
9.2 Fishery Cooperatives or Associations	12
9.3 Labor Organizations	12
9.4 Foreign Investment in the Domestic Fishery	12
10.0 DESCRIPTION OF SOCIAL AND CULTURAL FRAMEWORK OF DOMESTIC FISHERMEN AND THEIR COMMUNITIES.	12
11.0 DETERMINATION OF OPTIMUM YIELD	12
11.1 Specific Management Objectives	12
11.2 Specific Management Measures	14
12.0 A BIOLOGICAL AND FISHERIES PROFILE OF SPOTTED SEATROUT, <u>Cynoscion nebulosus</u>	15
12.1 Identity	15
12.1.1 Nomenclature	15
12.1.2 Taxonomy	15
12.1.2 Morphology	17
12.2 Distribution	18
12.2.1 General Distribution	18
12.2.2 Differential Distribution	18
12.2.2.1 Spawn, Larvae, and Juveniles	18
12.2.2.2 Adults	20

	Page
12.2.3 Determinants of Distribution	21
12.3 Life History	22
12.3.1 Reproduction	22
12.3.2 Pre-adult Phase	27
12.3.3 Adult Phase	27
12.3.4 Nutrition and Growth	27
12.3.5 Behavior	31
12.3.6 Contaminants	31
12.4 Population	34
12.4.1 Structure	34
12.4.2 Abundance, Density, Mortality, and Dynamics	40
12.4.3 Community Ecology	47
12.5 Exploitation	49
12.5.1 Commercial Exploitation	49
12.5.1.1 Fishing Equipment	49
12.5.1.2 Areas Fished	52
12.5.1.3 Fishing Seasons	52
12.5.1.4 Fishing Operations and Results. .	52
12.5.1.5 Incidental Catches	54
12.5.2 Recreational Exploitation	54
12.5.2.1 Fishing Equipment	54
12.5.2.2 Areas Fished	55
12.5.2.3 Fishing Seasons	57
12.5.2.4 Fishing Operations and Results. .	57

	Page
12.6 Social and Economic Implications	57
12.6.1 Values	57
12.6.2 Employment	59
12.6.3 Participation	59
12.6.4 Processors and Product Forms.	61
12.6.5 Import/Export	61
12.6.6 Gear Conflicts.	61
12.6.7 Commercial-Recreational Conflicts	61
12.7 Management and Protection	63
12.7.1 Regulatory Measures	63
12.7.2 Habitat Protection	71
12.7.3 Stocking	76
12.8 Current Research	76
12.9 Identification of Problems.	77
13.0 ACKNOWLEDGMENTS	78
14.0 REFERENCES	79
15.0 APPENDIX	99
15.1 Listing of members of Sciaenid Technical Committee, Sciaenid Board, and South Atlantic State/Federal Fisheries Management Board	99

3.0 INTRODUCTION

3.1 Development of the Plan

This spotted seatrout management plan was prepared through the Atlantic States Marine Fisheries Commission's (ASMFC) Interstate Fisheries Management Program under a contract between the North Carolina Division of Marine Fisheries and the National Marine Fisheries Service Southeast Region. The first phase in the development of this plan was the preparation of a profile summarizing available information on the biology of and fisheries for spotted seatrout (Mercer 1984, Section 12.0). This fishery management plan constitutes the second phase. The following six states participated in the development of the plan: Maryland, Virginia, North Carolina, South Carolina, Georgia, and Florida. The South Atlantic State-Federal Board, Sciaenid Board, and Sciaenid Technical Committee provided general guidance, policy, and technical expertise in the development of the plan.

3.2 Problems Addressed by the Plan

The status of spotted seatrout populations on the Atlantic Coast of the United States is unknown. Yield per recruit (YPR) analyses are being prepared for spotted seatrout populations in Florida; however, growth and mortality data are not available for other populations on the Atlantic Coast.

The population structure of spotted seatrout on the Atlantic Coast is not known. In the Gulf of Mexico tagging and enzyme electrophoretic studies have identified numerous subpopulations, suggesting that each population subunit could be treated as a separate management entity.

Annual landings of spotted seatrout are highly variable and fluctuations in catch seem to parallel the climatic conditions of the preceding winter(s). Large mortalities of spotted seatrout in shallow estuarine areas have been reported following severe cold spells. Other factors which may affect spotted seatrout landings include increased fishing effort and habitat alterations which may negatively affect the suitability of the estuarine habitat for spotted seatrout and thereby reduce the natural production of this species.

The spotted seatrout supports valuable commercial and recreational fisheries throughout much of its range. Increasing conflicts between the user groups have developed in recent years, particularly in the Gulf of Mexico. The Texas State Legislature resolved a spotted seatrout and red drum user conflict by creating game fish status for both species and thus eliminated the sale of these species by fishermen. This action has stimulated recreational fishermen in other states to pursue the enactment of similar restrictive laws and regulations.

Fisheries management decisions need to be based on an accurate knowledge of the biology, ecology, and population dynamics of spotted seatrout, and reliable catch and effort data for both the commercial and recreational fisheries. This management plan identifies spotted seatrout research and data needs and recommends management measures,

based on best available information, to perpetuate the spotted seatrout resource in fishable abundance throughout its range and generate the greatest possible economic and social benefits from its harvest and utilization over time.

4.0 DESCRIPTION OF THE STOCK

4.1 Species Distribution

Spotted seatrout, Cynoscion nebulosus, occur in estuarine and coastal waters from Cape Cod, Massachusetts to Carmen Island in the lower Gulf of Campeche, Mexico. Spotted seatrout are rare in and north of Delaware Bay, and the center of abundance extends from Florida to Texas.

The spotted seatrout is primarily an estuarine species associated with areas of submerged vegetation, sandy bottoms, and shell reefs. Spotted seatrout are year-round residents of estuaries along the South Atlantic coast, moving into deeper channels and holes, and occasionally offshore along the beaches to avoid extreme cold. Spotted seatrout are apparently migratory in Chesapeake Bay, moving offshore and south in fall and returning to the bay in spring. In North Carolina spotted seatrout are caught year-round in the estuaries, as well as in coastal oceanic waters in winter.

Differences in age and growth characteristics throughout the range, the non-migratory nature of spotted seatrout and the isolation of estuarine areas, and electrophoretic studies indicate that there are distinct subpopulations of spotted seatrout along the east coast of Florida and in the Gulf of Mexico. A Georgia tagging study indicated that movement was seasonal and generally short range. The stock structure on the Atlantic coast has not yet been determined.

4.2 Abundance and Present Condition

There are no data on relative abundance and present condition of the spotted seatrout population on the Atlantic coast. Commercial landings have fluctuated widely and have declined since 1950. Florida and North Carolina have accounted for the majority of Atlantic coast landings. Winter cold kills, environmental degradation, and fishing pressure have been cited as causes of declines of spotted seatrout landings.

4.3 Ecological Relationships

The information presented in this section is summarized from the profile (Section 12.0).

Reproduction - Spotted seatrout mature between one and three years of age, and males tend to mature at a smaller size (250 mm TL) than females (290 mm TL). The overall sex ratio was about 2:1 (females to males) in most studies. Spawning occurs at night, from April through September, both within estuaries and in the near-shore coastal waters. Estimates of spotted seatrout fecundity range from 1.4×10^4 to 1.6×10^7 eggs. Young-of-the-year are generally associated with seagrass beds in areas where these beds occur.

Age and Growth - The maximum age reported for spotted seatrout is 15 years in Chesapeake Bay. Eight and 10 years were the maximum ages reported for spotted seatrout in Georgia and Florida, respectively. Females attain a greater maximum age than males throughout the range. Mean back-calculated lengths at age and von Bertalanffy growth parameters are presented in Tables 12-6 and 12-7.

Food and Feeding - Spotted seatrout are carnivorous, feeding primarily on crustaceans (penaeid shrimp and crabs) and fishes (anchovies, menhaden, mullet, pinfish, and silversides). Juveniles (<150 mm SL) feed on copepods, mysids, caridean and palaemonid shrimps, amphipods, and polychaetes (Table 12-5).

Competitors and Predators - Klima and Tabb (1959) listed red drum, spot, croaker, silver perch, grouper, and sea catfish as competitors with spotted seatrout in Florida waters. Snook, gafftopsail catfish, common jack, mangrove snapper, and spotted seatrout were listed as predators.

Parasites, Diseases, Injuries and Abnormalities - Overstreet (1983) presented a partial list of parasites of spotted seatrout and reviewed the literature on diseases, mortalities, and abnormal conditions found in spotted seatrout.

4.4 Estimate of Maximum Sustainable Yield

Yield per recruit models were generated for the Everglades National Park recreational fishery for spotted seatrout (Rutherford 1982). Condrey et al. (1984) calculated yield per recruit for spotted seatrout in U.S. Gulf of Mexico estuaries based on age, growth, and mortality data presented in the literature. Preliminary estimates of yield per recruit for spotted seatrout populations on the east and west coasts of Florida were presented at a recent stock assessment workshop.

4.5 Probable Future Condition

The future condition of spotted seatrout populations on the Atlantic coast is dependent on larval recruitment, maintaining or improving environmental quality, and management efforts to conserve the resource.

5.0 DESCRIPTION OF HABITAT

5.1 Condition of the Habitat

Estuarine habitats, where nearly the entire commercial and recreational catches of spotted seatrout are made, have deteriorated rapidly since approximately 1950, mostly as a result of industrial and human population growth. The National Estuary Study, completed in 1970, indicated that 73% of the Nation's estuaries had been moderately or

¹Murphy, M. 1984. Red drum and spotted seatrout stock assessment in Florida. Paper present at Florida Department of Natural Resources Stock Assessment Workshop, November 16 and 17, 1984, St. Petersburg, Florida.

severely degraded (Gusey 1978, 1981). Damage and/or destruction of estuaries has largely been by dredging and filling for waterfront property, dredging of navigation channels, construction of causeways and bridges, installation of ports and marinas, alteration of freshwater flow, and pollution. Unfortunately the effects of habitat alterations have rarely been quantified.

5.2 Habitat Areas of Particular Concern

The association of juvenile and adult spotted seatrout with seagrass beds is well documented (Pearson 1929; Miles 1950; Moody 1950; Reid 1954; Tabb 1958). Seagrass beds along the coast of Mississippi were virtually destroyed during Hurricane Camille in August 1969 (Lorio and Perret 1980); however, it is not known what effects this had on spotted seatrout populations in Mississippi Sound. Declines in spotted seatrout landings in Tampa Bay, Florida have occurred concomitantly with large declines in seagrass acreage (Taylor and Salomar 1968; Lewis 1977; Lewis and Phillips 1980). An unprecedented decline in submerged aquatic vegetation has occurred in Chesapeake Bay in the last 15 to 20 years (Orth and Moore 1983). Major changes in vegetation patterns began in 1972, the year of Tropical Storm Agnes, which lowered salinities for periods of up to four weeks and transported large quantities of suspended sediment into the estuarine system. The causes of the Chesapeake Bay decline in seagrass beds are not known but may be related to nutrient enrichment affecting the quantity and quality of light reaching the plant surface. Implications for species inhabiting grass beds have not been determined but could be considerable.

5.3 Habitat Protection Programs

State Programs

The coastal states have enacted coastal zone management laws to regulate shoreline development and protect the coastal zone. State habitat protection regulations are summarized in Table 12-23.

Federal Programs

The Coastal Zone Management Act of 1972, 16 USC 1451

The Act establishes a national policy and initiates a national program to encourage state planning for the management, beneficial use, protection and development of the Nation's coastal zones (generally, the submerged lands and waters of the territorial sea and the adjacent shorelands having a direct and significant impact on such waters).

Fish and Wildlife Coordination Act of 1956, USC 742(a)-754

Established a comprehensive national policy on fish and wildlife resources; authorized programs and investigations that may be required for the development, advancement, management, conservation and protection of the fisheries resources of the United States.

National Environmental Policy Act of 1969, 42 USC 4321-4347

Requires detailed environmental impact statements of proposals for legislation and other major Federal actions which may significantly affect the quality of the human environment. Prior to making the detailed statement, the responsible Federal official is required to consult with and obtain the comments of any Federal agency which has jurisdiction by law or special expertise with respect to any environmental impact involved. Also requires that documents must be available to the public and their comments must be considered.

The Ports and Waterways Safety Act of 1972, 33 USSC 1221-1227

This Act deals with transportation and pollution problems resulting from operation and casualties of vessels carrying oil and other hazardous substances. It is designed to protect coastal waters, living resources, recreational resources and scenic values.

Federal Water Pollution Control Act, and Amendments of 1972, 33 USC 1251-1376

This Act initiated major changes in the enforcement mechanism of the Federal water pollution control program from water quality standards to effluent limits. Among other things, it requires that permits be issued by the Environmental Protection Agency or the States for discharge of effluents into waters of the United States.

The Marine Protection, Research and Sanctuaries Act of 1972 (The Ocean Dumping Act), 33 USC 1401-1444

This Act regulates the transportation from the United States of material for dumping into the oceans, coastal and other waters, and the dumping of material from any source into waters over which the United States has jurisdiction. The Environmental Protection Agency is empowered to issue permits for transportation or dumping where it will not unreasonably degrade or endanger human health, welfare or amenities, or the marine environment, ecological systems or economic potentialities. Section 106 of the Act provides for the provisions of the Fish and Wildlife Coordination Act to apply.

Marine Mammals Protection Act of 1972, 16 USC 1361-1407

This Act, with certain exceptions, places a moratorium on the taking and importation of all marine mammals and marine mammal products. It makes the Secretary of Commerce responsible for protecting whales, porpoises, seals, sea lions; and the Secretary of the Interior responsible for all other marine mammals, specifically sea otters, walruses, polar bears and manatees. Also protects the habitat of marine mammals, including food sources.

Endangered Species Act of 1974, PL 93-205

This Act gives the Departments of Commerce and Interior regulatory and statutory authority on endangered and threatened fauna and flora not included in previous Acts. The purpose of the Act is to conserve endangered and threatened species and the ecosystems upon which they depend.

Deepwater Port Act of 1974, 33 USC 1501-1524

Establishes procedures for the location, construction and operation of deepwater ports off the coasts of the United States.

Magnuson Fishery Conservation and Management Act of 1976

Establishes a fishery conservation and management regime to be implemented by the Secretary of Commerce. Establishes a fishery conservation zone extending from the limits of the territorial sea to 200 nautical miles from the baseline from which the territorial sea is measured. The Act defines fishery resource to include ". . . any habitat of fish", and enjoins the Secretary to carry out a research program which must include ". . . the impact of pollution on fish, the impact of wetland and estuarine degradation, and other matters. . ."

National Ocean Pollution Research and Development and Monitoring Planning Act of 1978, PL 95-273

Designates NOAA as the lead agency in the development of a comprehensive five-year plan for a Federal program relating to ocean pollution research, development and monitoring. This plan is to provide for the coordination of existing Federal programs relating to the oceans and for the dissemination of information emerging from these programs to interested parties. In addition, the plan shall provide for the development of a base of information necessary to the utilization, development and conservation of ocean and coastal resources in a rational, efficient and equitable manner.

6.0 FISHERY MANAGEMENT JURISDICTION, LAWS, AND POLICIES

6.1 Management Institutions

The U.S. Department of Commerce, acting through the Fishery Management Councils, pursuant to P.L. 94-265 (Magnuson Fishery Conservation and Management Act), has authority to manage stocks throughout the range that are harvested predominantly in the FCZ.

6.2 Treaties and International Agreements

Foreign fishing is regulated by P.L. 94-265 pursuant to which Governing International Fishing Agreements are negotiated with foreign nations for fishing within the FCZ.

6.3 Federal Laws, Regulations and Policies

The only known federal law that can regulate the management of the spotted seatrout fishery is P.L. 94-265.

6.4 State Laws, Regulations and Policies

The fisheries for spotted seatrout have been conducted almost entirely within the internal waters of the states and in the territorial sea which extends 5.6 km (3 n mi) offshore on the Atlantic coast. Therefore, management has been by individual state regulation. Regulations and methods of promulgating them vary among states and are summarized in Table 12-18. The only regulations specifically dealing with spotted seatrout are minimum size limits of 23 cm (9 in) TL in Maryland and 30 cm (12 in) FL in Florida.

6.5 Local and Other Applicable Laws, Regulations, and Policies

No local or other laws, regulations, or, policies are known to exist relative to the spotted seatrout fishery.

7.0 DESCRIPTION OF FISHING ACTIVITIES

7.1 History of Exploitation

Spotted seatrout are fished both commercially and recreationally. Commercial landings of spotted seatrout have been recorded for the Gulf of Mexico since 1887 and for the Atlantic since 1923. Gulf of Mexico landings have generally accounted for 60-80% of total United States spotted seatrout commercial landings.

Commercial landings of spotted seatrout have been reported from Maryland to Florida on the Atlantic coast (Table 7-1). Total Atlantic coast landings were highest in 1945 (2,495 mt) and have since declined, fluctuating between 200 and 600 mt over the last two decades. Florida and North Carolina account for most of the Atlantic coast spotted seatrout landings. Florida landings peaked at 1,985 mt in 1945 and declined to 200-400 mt from 1957 to the present. North Carolina landings have fluctuated with peak landings in 1932 (850 mt) and a low of 38 mt in 1982. Maryland landings of spotted seatrout never exceeded 12 mt and none have been reported since 1956. Virginia landings were highest in 1944 (345 mt) and fluctuated between 1 and 89 mt from 1947 to 1983. Spotted seatrout landings in South Carolina peaked at 101 mt in 1956 and declined to <1-4 mt from 1976 to the present. Georgia landings were highest in 1936 at 52 mt and ranged from <1 to 20 mt from 1950 to the present.

Recreational fishing surveys indicate that the sport fishery catch of spotted seatrout probably exceeds the commercial harvest (Tables 7-1 and 7-2). The total estimated recreational catches of spotted seatrout on the Atlantic coast for 1965 (8,267 mt) and 1970 (11,368 mt) exceeded the commercial landings for those years (426 and 545 mt, respectively) by factors of 19 and 21. The recreational surveys for 1979-1980, which are believed to be more reliable than previous surveys, indicate that the

recreational catch was from three to 11 times greater than the commercial catch for those years. The South Atlantic subregion (North Carolina-Florida) accounts for most of the recreational catch of spotted seatrout.

7.2 Domestic Commercial and Recreational Fishery Activity

Commercial Fishery

The commercial fishery for spotted seatrout is primarily a mixed species fishery conducted within estuarine and nearshore coastal waters. The principal commercial methods used to harvest spotted seatrout include various types of gill nets, haul seines, pound nets, hand lines, troll and trot lines, trammel nets, and otter trawls.

Fishing gear varies among states and is partly a function of gear efficiency in different areas as well as a function of state or local laws (Table 12-12). In Virginia the majority of spotted seatrout are landed in common haul seines, with pound nets, gill nets, and fish trawls contributing the remainder. North Carolina landings are mainly derived from common and long haul seines and gill nets, with smaller quantities from pound nets, hand lines and otter trawls (fish, crab and shrimp). In South Carolina spotted seatrout are caught by gill nets, shrimp trawls, and hand lines. Georgia landings are chiefly by hand lines. In Florida spotted seatrout are primarily caught in runaround gill nets, as well as hand lines, common haul seines, trammel nets, troll lines, and shrimp trawls.

Recreational Fishery

Spotted seatrout are caught by hook and line in estuaries, inlets, and the surf. Results of the marine recreational fishery statistics surveys, 1965-1980, indicate that 69-97% of the spotted seatrout were caught in inland or estuarine waters (Table 7-3). The principal mode of fishing was private/rental boats (Table 7-3).

7.3 Foreign Fishing Activities

There is no foreign catch of spotted seatrout because spotted seatrout generally do not occur greater than 3 mi from shore.

8.0 DESCRIPTION OF ECONOMIC CHARACTERISTICS OF THE FISHERY

8.1 Domestic Harvesting Sector

Spotted seatrout contributed more to the total value of U.S. sciaenid landings between 1960 and 1974 than any other species (Cato 1981). Total value of spotted seatrout landings have ranked third behind croaker and weakfish since 1977. Atlantic coast values of spotted seatrout landings have fluctuated, and have increased since 1979 (Figure 12-4). The real (deflated) price of spotted seatrout declined from 1967 to 1977 on the Atlantic coast (Table 12-16).

Table 7-1. Atlantic coast commercial landings of spotted seatrout (metric tons), 1950-1983.

Year	Maryland	Virginia	North Carolina	South Carolina	Georgia	East Coast Florida	Atlantic Coast Total
1950	-	42	219	12	15	661	949
1951	-	29	121	21	14	568	753
1952	-	33	292	39	11	675	1,053
1953	-	39	287	23	8	252	609
1954	-	47	365	12	9	556	989
1955	-	47	201	63	20	413	744
1956	*	89	177	101	5	485	857
1957	-	54	262	25	2	402	745
1958	-	27	80	9	1	346	463
1959	-	63	176	17	*	348	504
1960	-	25	78	13	*	404	520
1961	-	33	05	25	1	340	494
1962	-	13	93	12	*	343	461
1963	-	12	105	22	2	363	504
1964	-	11	93	27	1	347	479
1965	-	18	79	16	4	309	426
1966	-	5	53	11	1	329	398
1967	-	2	56	1	3	282	334
1968	-	3	44	5	1	289	342
1969	-	8	86	4	1	308	407
1970	-	30	183	4	5	323	545
1971	-	20	153	11	7	222	413
1972	-	6	228	8	12	288	542
1973	-	4	288	3	12	302	598
1974	-	12	304	4	7	299	626
1975	-	33	287	8	14	243	585
1976	-	18	289	3	14	241	565
1977	-	2	147	*	7	224	380
1978	-	3	44	*	1	183	231
1979	-	2	48	1	2	216	269
1980	-	1	78	4	2	253	338
1981	-	2	51	*	*	334	387
1982	-	2	38	1	2	330	373
1983	-	2	75	2	3	206	288

- None reported

* Less than 1 mt

Table 7-2. Estimated weight of spotted seatrout caught by marine recreational fishermen by state and subregion for the Atlantic coast (kg x 1000), 1965-1980.

<u>Area</u>	<u>Year</u>			
	1965 ¹	1970 ²	1979 ³	1980 ³
Mid-Atlantic				
Delaware	*	*	20	*
Maryland	*	*	109	*
Virginia	*	*	431	*
Subregion Total	*	*	560	*
South Atlantic				
North Carolina	*	*	8	112
South Carolina	*	*	26	249
Georgia	*	*	351	211
Florida E. Coast	*	*	2,005	409
Subregion Total	8,267	11,368	2,390	981
TOTAL	8,267	11,368	2,950	981

*None reported

²Deuel 1973

¹Deuel and Clark 1968

³Anonymous 1984

Table 7-3. Estimated total number of spotted seatrout caught by marine recreational fishermen by area and mode of fishing, 1965-1980, for the Atlantic coast.

Survey year	Region	Area of Fishing		Mode of fishing				Beach/ Bank
		Ocean	Inland	Private/ Rental	Party/ Charter	Man- Made		
----- T H O U S A N D S -----								
1965 ¹	South Atlantic	2,343	10,216	10,173	180	1,412		794
1970 ²	South Atlantic	4,374	9,618	7,686	2,187	3,433		686
1979 ³	Mid-Atlantic	-	393	401	-	-		*
	South Atlantic	85	3,282	3,151	*	92		124
1980 ³	South Atlantic	72	1,675	1,682	12	193		91
*None reported		¹ Deuel and Clark 1968		³ Anonymous 1984				
-<30,000 reported		² Deuel 1973						

8.2 Domestic Processing Sector

Spotted seatrout is almost strictly a fresh fish product, sold to either local wholesale or retail markets.

8.3 International Trade

Imports of spotted seatrout from Mexico are substantial (Perret et al. 1980) (Table 12-17). There are no records of exports of spotted seatrout from the U.S.

9.0 DESCRIPTION OF THE BUSINESSES, MARKETS AND ORGANIZATIONS ASSOCIATED WITH THE FISHERY

9.1 Relationship Among Harvesting and Processing Sectors

Spotted seatrout are sold as fresh in-the-round or gutted. A small percentage is sold as frozen and gutted or as fresh or frozen fillets (Perret et al. 1980).

9.2 Fishing Cooperatives or Associations

No data are available on fishing cooperatives or associations.

9.3 Labor Organizations

No data are available on labor organizations.

9.4 Foreign Investment in the Domestic Fishery

No information is available on foreign investment in the domestic fishery.

10.0 DESCRIPTION OF SOCIAL AND CULTURAL FRAMEWORK OF DOMESTIC FISHERMEN AND THEIR COMMUNITIES

Uniform socio-economic data on fishing communities are not available.

11.0 DETERMINATION OF OPTIMUM YIELD

The goal of this plan is to perpetuate the spotted seatrout resource in fishable abundance throughout its range and generate the greatest possible economic and social benefits from its harvest and utilization over time.

11.1 Specific Management Objectives

The following objectives have been adopted for achievement of the goal:

1. Attain over time optimum yield.

2. Maintain a spawning stock sufficient to minimize the possibility of recruitment failure.

Tagging, age and growth, and electrophoretic studies have indicated the existence of nearly independent groups of spotted seatrout throughout much of the range. Therefore, there may be relatively little recruitment from outside or adjacent estuaries. Although adapted to the shallow, highly variable estuarine waters, spotted seatrout are very sensitive to rapid changes in temperature. Winter-time cold shock of juveniles and adults has been cited as a primary factor in local and coastwide declines in spotted seatrout. Adverse environmental conditions and/or damage to the habitat, in combination with increasing fishing pressure, could affect production of localized populations of spotted seatrout.

3. Promote conservation of the stocks in order to reduce interannual variation in availability and increase yield per recruit.

The presence in the population of fish of several ages tends to prevent sudden fluctuations in the catch resulting from the poor survival of fish of one particular year class. Management measures such as minimum size limits will conserve a year class and increase yield per recruit.

4. Promote the collection of economic, social, and biological data required to effectively monitor and assess management efforts relative to the overall goal.

Effective management of the spotted seatrout resource is hindered by the lack of reliable and consistent coastwide catch and effort data from the commercial and recreational fisheries. There is a lack of information on population dynamics, including stock size, age and size composition, and natural and fishing mortality rates, required for the development of yield models. Very little is known about the economics of the commercial fishery or the economic benefits from recreational fishing. There is a lack of sociological information on recreational and commercial fishermen.

5. Promote research that improves understanding of the biology and fisheries of spotted seatrout.

Life history and population data throughout the range are required to effectively manage the fisheries. Existing evidence indicates that separate subpopulations of spotted seatrout occur in estuaries along the coasts of Florida and the Gulf of Mexico. The population structure on the Atlantic coast has not been determined; however, if local populations do exist, life history parameters need to be defined for each population. Other research needs include determining habitat requirements and the effects of environmental factors on stock size.

6. Promote harmonious use of the resource among various components of the fishery through the coordination of management efforts among the various political entities having jurisdiction over the spotted seatrout resource.

Spotted seatrout on the Atlantic Coast are harvested commercially primarily in mixed-species fisheries and recreationally. As with any species in demand by more than one user group, recreational-commercial fishery conflicts have developed. Management decisions need to be based on all of the necessary biological information, as well as social, economic, and political factors.

7. Promote determination and adoption of standards of environmental quality and provide habitat protection necessary for the maximum natural production of spotted seatrout.

The estuarine system as a whole must be considered when dealing with spotted seatrout management. The shallows and grassbeds have been identified as critically important nursery areas for spotted seatrout. Habitat alterations, pesticides, and other pollutants entering estuaries may affect the production of spotted seatrout.

11.2 Specific Management Measures

The following management strategy is recommended to obtain the above objectives:

1. A minimum size limit of 12 in TL with comparable mesh size regulations in directed fisheries for spotted seatrout. A directed fishery is defined as containing at least 60% spotted seatrout by weight.

The purpose of a minimum size limit is to protect the spawning stock and increase yield per recruit. Present minimum size limits for spotted seatrout are 12 in FL in Florida and 9 in TL in Maryland. The recommended minimum size is the length at which 50% of the females reach maturity.

2. Collection of improved catch and effort data from the commercial and recreational fisheries, including size composition of the catch and social and economic data.
3. Additional measures, not specifically recommended in this plan, such as creel limits, catch quotas, limited entry, area closures, and gear restrictions, may be implemented in the future.
4. The following monitoring and research activities are recommended:
 - a. stock identification,
 - b. mortality estimates and yield modeling,
 - c. habitat requirements,
 - d. effects of environmental factors on stock size,
 - e. development of a prerecruit index,
 - f. mesh size selectivity, and
 - g. social and economic analyses.

12.0 A BIOLOGICAL AND FISHERIES PROFILE OF SPOTTED SEATROUT, Cynoscion nebulosus

12.1 Identity

12.1.1 Nomenclature

The valid name for spotted seatrout is Cynoscion nebulosus (Cuvier) 1830 (Figure 12-1). The following synonymy is after Jordan and Evermann (1896):

Labrus squeteague var. maculatus, Mitchill, 1815
Otolithus nebulosus, Cuvier and Valenciennes, 1830
Otolithus carolinensis, Cuvier and Valenciennes, 1833
Otolithus drummondi, Richardson, 1836
Cestreus carolinensis, Gronow, 1854
Cynoscion carolinensis, Jordan and Gilbert, 1878
Cynoscion maculatum, Jordan and Gilbert, 1882
Cestreus nebulosus, Jordan and Eigenmann, 1889

12.1.2 Taxonomy

Classification follows Greenwood et al. (1966). Taxa higher than superorder are not included.

Superorder: Acanthopterygii
 Order: Perciformes
 Suborder: Percoidei
 Family: Sciaenidae
 Genus: Cynoscion
 Species: Cynoscion nebulosus

The spotted seatrout is one of 33 members of the family Sciaenidae found along the Atlantic, Gulf and Pacific coasts of the United States (Robins et al. 1980). Members of this family are commonly known as drum fishes or croakers because of the drumming or croaking sounds they produce by vibrating their swimbladders (Bigelow and Schroeder 1953). Chao (1978) assessed the phylogenetic relationships of western Atlantic Sciaenidae on the basis of swimbladder, otoliths, and external morphology. He also presented a field key to the genera and species, including meristics and species ranges. Hildebrand and Cable (1934) provided a key to eggs and young (<35 mm TL) of Atlantic coast drums.

There are three other members of the genus Cynoscion found along the Atlantic and Gulf coasts of the United States: weakfish, C. regalis; silver seatrout, C. nothus; and sand seatrout, C. arenarius. Mohsin (1973) compared the osteology of these four species and hypothesized that there are two phyletic lines within the genus Cynoscion: one comprised of C. nebulosus and C. arenarius and the other of C. nothus and C. regalis. Results of an electrophoretic investigation refuted Mohsin's conclusion and indicated that C. arenarius should be regarded as a subspecies of C. regalis (Weinstein and Yerger 1976a). These




Figure 12-1. Spotted seatrout, *Cynoscion nebulosus* (Cuvier), 1830, illustrated by Goode, 1884.

results supported conclusions of previous morphological (Ginsburg 1929) and ecological (Tabb 1966) studies that C. nebulosus is the most divergent of the four forms. An electrophoretic study of C. nebulosus from various estuaries along the Atlantic and Gulf coasts indicated that each estuary may contain a discrete subpopulation. (Weinstein and Yerger 1976b).

Spotted seatrout is the common name given Cynoscion nebulosus by the American Fisheries Society (Robins et al. 1980). Other common names are spotted weakfish, spotted squeteague, salmon trout, simon trout, spotted trout, speckles, specs, speckled trout, trout, seatrout, black trout, salmon, southern squeteague and winter trout (Smith 1907; Hildebrand and Schroeder 1928; Shiino 1976).

12.1.3 Morphology

The following description is that of Johnson (1978), summarized from Jordan and Evermann (1896), Welsh and Breder (1923), Hildebrand and Schroeder (1928), Hildebrand and Cable (1934), Miller and Jorgenson (1973), and Chao (1976).

D. X (rarely IX or XI)-I, 24-28, A. II, 9-12 (typically 10-11); C. 9+8, procurrent rays 6-9+5-7; V. I, 5; lateral line scales 90-102, scales between anal fin origin and lateral line 11-12; vertebrae 13+12; gill rakers 6-9 on lower limb; branchiostegals 7; a pair of large canine-like teeth at tip of upper jaw; remaining teeth small conical, set in narrow bands with outer row slightly enlarged in upper jaw and inner row distinctly enlarged in jaw; no teeth on vomer, palatines, or tongue.

Head 2.9-3.5, depth 3.4-4.5 in SL; snout 3.7-4.2, eye 4.4-5.3, interorbital 4.5-5.9, maxillary 2.2-2.3, pelvic fin 1.8-2.2 in head.

Body elongate and somewhat compressed; back a little elevated; head long; snout pointed; mouth large, oblique; lower jaw projecting; maxillary reaching to or nearly to posterior margin of eye. Scales moderate, thin, all ctenoid, fins scaleless, except for 1-10 rows of small scales at dorsal and anal fin bases. Dorsal fin continuous or slightly separate, the spines weak, flexible; anal fin small, second spine very weak, caudal fin straight to somewhat emarginate. Preopercular margin smooth, sometimes ciliated, never with strong serrations.

Pigmentation: Color dark gray above, with sky blue reflections, shading to a silvery below; upper parts of sides with numerous round black spots extending onto dorsal and caudal fins; fins pale to yellowish green.

Readily distinguished from related species by the round black spots on upper parts of body and on dorsal and caudal fins, the small scales, and the scaleless median fins.

Development of body proportions and meristic characters of larvae and juveniles (1.9-32.2 mm NL and SL) were reported by Powles and Stender (1978). Standard length-total length relationships for spotted seatrout in Georgia (Jorgenson and Miller 1968), Louisiana (Hein et al. 1980), Mississippi (Overstreet 1983), and Texas (Harrington et al. 1979) are presented in Table 12-1.

12.2 Distribution

12.2.1 General distribution

Spotted seatrout occur along the Atlantic and Gulf coasts of the United States from Cape Cod, Massachusetts to Carmen Island in the lower Gulf of Campeche, Mexico (Welsh and Breder 1923; Mather 1952; Tabb 1966; Yanez-Arancibia et al. 1980). They are rare in and north of Delaware Bay (Welsh and Breder 1923), and the center of abundance is from Florida to Texas (Pearson 1929).

12.2.2 Differential Distribution

12.2.2.1 Spawn, Larvae, and Juveniles

Spotted seatrout eggs have not been identified in field collections and data on the preferred spawning habitat are conflicting. Information on spawning areas in the Gulf of Mexico was summarized by Hein and Shepard (1979a) and Perret et al. (1980). The distribution of larvae and gravid and spent adults indicates that spawning occurs both within estuaries and in near-shore coastal waters along the Atlantic and Gulf coasts (Pearson 1929; Miles 1950, 1951; Stewart 1961; Tabb and Manning 1961; Tabb 1966; Jannke 1971; King 1971; Christmas and Waller 1973; Mahood 1975; Powles and Stender 1978; Houde et al. 1979; Brown 1981; Overstreet 1983).

Spotted seatrout larvae and postlarvae have been collected in channels and passes entering Florida, Louisiana, and Texas bays (Jannke 1971; King 1971; Sabins and Truesdale 1974; Robison in press) and in the eastern Gulf of Mexico within the 15 m isobath (Houde et al. 1979). Miles (1950) reported a close association between postlarvae and widgeon grass (*Ruppia maritima*) in Texas bays. Hildebrand and Cable (1934) collected only 17 spotted seatrout larvae (<5.0 mm) from 11 km offshore to within the Newport River estuary in several years of intensive sampling in the vicinity of Beaufort, North Carolina. All larger specimens were caught in the estuary. Williams and Deubler (1968) collected six metamorphosing spotted seatrout (16.0 mm TL) during two years of sampling in the lower Neuse and Pamlico rivers, N.C. In South Carolina larvae and postlarvae (1.9-32.2 mm SL) were collected in lower portions of the estuaries and in tidal passes (Powles and Stender 1978). Setzler (1977) sampled along a transect from 10.5 km offshore to the head of Doboy Sound, Sapelo Island, Georgia and found that spotted seatrout postlarvae were restricted to the estuary. In the Indian River lagoon system of east-central Florida young (<20 mm) were observed in moderately deep water (<3 m) over algae and muddy sand bottom (Tabb 1961). Extensive sampling there on the shallow flats failed to produce postlarvae or juveniles. The low number of specimens collected along

Table 12-1. Standard length-total length relationships for spotted seatrout as reported in the literature.

Reference	Location	Size Range (mm TL)	N	Sex	Relationship	r
Jorgenson and Miller (1968)	Georgia	10-72	20	-	TL = 1.020 + 1.224 SL SL = -0.764 + 0.815 TL	- -
Harrington et al. (1979)	Texas	53-858	9857	-	TL = 11.804 + 1.138 SL	0.997
Hein et al. (1980)	Louisiana	21-629	1208	-	SL = -3.883 + 0.865 TL	0.995
Overstreet (1983)	Mississippi	55-624	1680	females	TL = 10.659 + 1.128 SL SL = -8.354 + 0.883 TL	0.998 0.998
		55-535	1089	males	TL = 10.165 + 1.129 SL SL = -7.465 + 0.880 TL	0.997 0.997

the Atlantic coast prevents any conclusion as to the preferred habitat of larval spotted seatrout.

Young-of-the-year spotted seatrout are generally associated with seagrass beds in estuaries. Pearson (1929) collected hundreds of juveniles (20-30 mm TL) along the grassy shore lines of remote Texas bays, small restricted bayous, and creeks. In Chesapeake Bay, Virginia, juvenile spotted seatrout (24-140 mm TL) were collected in seagrass beds from July to October (Orth and Heck 1980; Brown 1981). In November young-of-the-year were caught by trawl in the channels (18-23 m) of the York and James rivers (Brown 1981). In North Carolina juveniles were collected from April to September, mainly in areas of seagrass (Spitsbergen and Wolff 1974; Wolff 1976; Purvis 1976; Miller and Dunn 1980). A few juveniles were collected in trawl surveys of South Carolina estuaries (Lunz and Schwartz 1969; Turner and Johnson 1972) and in seine and rotenone collections from an intertidal creek (Cain and Dean 1976). Spotted seatrout juveniles were trawled and seined in Georgia estuaries and along the beaches in the surf zone (Miller and Jorgenson 1969; Dahlberg 1972; Mahood 1975). More were usually taken in tidal pools and small creeks at low tide when they could not hide in the grasses along the banks (Mahood 1975). In the Indian River area of Florida, juvenile spotted seatrout were collected in areas of sand and seagrass (Tabb 1961; Jones et al. 1975). The smallest individuals (8-50 mm SL) were taken in moderately deep channels (<3 m) and appeared to disperse to shallower grassy bays with increasing size (Tabb 1961). Juveniles (13-173 mm FL) were collected over mud and sand bottom in low salinity areas (0-11.1 ppt) of the St. Johns River system, Florida (Tagatz 1967).

12.2.2.2 Adults

The spotted seatrout is primarily an estuarine species and is most abundant in the confines of semi-landlocked lagoons and quiet estuaries (Tabb 1958). Along the coasts of Florida, Alabama, Mississippi, and Texas, spotted seatrout are primarily found in large areas of shallow, quiet, brackish water with extensive submerged vegetation characterized by turtle grass (Thalassia testudinum), shoal grass (Halodule wrightii), and widgeon grass, with adjacent deep areas (3-6 m) used for refuge from high summer temperatures and winter cold (Pearson 1929; Miles 1950; Tabb 1958; Lorio and Perret 1980; Zieman 1982). They are also found in deeper bays and around oyster reefs along the Texas coast (Hoese and Moore 1977). In Louisiana spotted seatrout are associated with sandy bottoms, submerged or emergent islands, shell reefs, areas of submerged vegetation, areas where some type of structure exists (e.g., oil platforms), and deep bayous and canals (Lorio and Perret 1980). Gilmore (1977) reported that spotted seatrout are common on grassflats and sand bottom, occasional in mangroves and inlets in the Indian River lagoon, and rare in the offshore benthic-open shelf habitat in that region. The absence of seagrass beds along the coasts of South Carolina and Georgia apparently does not limit the distribution of spotted seatrout in these areas (Hoese 1973). In South Carolina spotted seatrout are usually found around shell banks in creeks, rivers, and sounds (Bearden 1961; Bearden and Farmer 1972; Hicks 1972; Shealy et al. 1974). In Georgia adults were found in all sectors of the estuaries and

along ocean beaches, usually at depths of 1 to 3 m (Dahlberg 1972; Mahood 1975). In Chesapeake Bay spotted seatrout tend to stay in shallow creeks and rivers adjacent to beds of eelgrass (*Zostera marina*) and widgeon grass, although they will move into deep holes during midsummer (Brown 1981).

Spotted seatrout are year-round residents of estuaries along the South Atlantic coast and in the Gulf of Mexico, moving into deeper channels and holes and occasionally offshore along the beaches to avoid extreme cold (Pearson 1929; Moody 1950; Simmons 1951; Guest and Gunter 1958; Tabb 1958, 1966; Dahlberg 1972; Mahood 1975; Hein and Shepard 1979b; Music and Pafford 1984). Spotted seatrout are found year-round in the sounds and mouths of rivers in North Carolina (Smith 1907; Hildebrand and Cable 1934; Roelofs 1951).

Spotted seatrout are apparently migratory in the northern portion of their range. Hildebrand and Schroeder (1928) reported that spotted seatrout were caught by seines in the lower part of Chesapeake Bay in early fall until cold weather arrived, at which time they left the bay and moved south. Analysis of Virginia Saltwater Fishing Tournament citation records revealed that the largest catches of spotted seatrout in Chesapeake Bay were made in May and October, corresponding to spring and fall migrations (Brown 1981). In November spotted seatrout are caught by sport fishermen in the deep channel areas of the Chesapeake Bay Bridge Tunnel, where they are often associated with weakfish. There is some indication that a portion of the North Carolina population may be migratory. Spotted seatrout are caught in trawls and gill nets off the coast of North Carolina from November through March. Whether these catches are composed of fish from northern areas or from North Carolina estuaries, or a mixture, is not known. An account of spotted seatrout habits in North Carolina in the 1800s stated that this species first made its appearance in February on its way from the south, remained in the vicinity of sounds and inlets until about May, gradually proceeded northward, and reappeared on the coast of North Carolina in September (Goode 1884).

12.2.3 Determinants of Distribution

Tabb (1958) listed the following ecological characteristics which appeared to be of greatest importance in determining the abundance and "success" of spotted seatrout in Florida: (1) large areas of shallow, quiet, brackish waters; (2) extensive grassy areas usually dominated by turtle grass and shoal grass; (3) areas of 3-6 m depth adjacent to grass flats to be used for refuge from winter cold; (4) an abundant food supply, viz., grazing crustaceans and suitable size fish; (5) absence of predators; (6) absence of competitors; and (7) suitable temperature range of 15-27°C. The association of both juvenile and adult spotted seatrout with seagrass beds, as well as other types of habitats, is well documented (Sections 12.2.2.1 and 12.2.2.2). Temperatures below 7-10°C cause spotted seatrout in Florida to move into ocean inlets or offshore along beaches for brief periods of time (Tabb 1958). Temperature was also determined to be a factor in Georgia estuaries, with movement of spotted seatrout into deeper waters at temperatures > 25°C or < 16°C (Mahood 1975).

The spotted seatrout is a euryhaline species, recorded from fresh water (0.2 ppt) (Perret 1971) to hypersaline conditions of 75 ppt (Simmons 1957). Loman (1978) reported that largest catches of spotted seatrout in Mississippi were caught between 20 and 35 ppt. The optimal salinity reported from laboratory studies is 20 ppt (range: 10-45 ppt) at 28°C based on standard routine and maximum sustained respiratory metabolic rates (Wakeman and Wohlschlag 1977; Wohlschlag and Wakeman 1978). Maximum sustained swimming speeds occurred at 20-25 ppt at 28°C, and swimming performance was reduced at salinities above or below this range. Tabb (1966) reported a normal salinity range of 5-30 ppt in Florida and that sudden changes in salinity, such as caused by tropical storms or hurricanes, may cause mass migrations or mortalities. Reported optimal salinity ranges for larvae were 20-35 ppt (Arnold et al. 1976) and 18.6-37.5 ppt (Taniguchi 1980).

There are no data relating the distribution of spotted seatrout to dissolved oxygen concentrations. Vetter (1977) reported from laboratory studies that the oxygen requirements for spotted seatrout at 28°C (a normally prevailing summer temperature in coastal Texas waters) with salinities of 10, 20, and 30 ppt were 210, 125 and 230 mg O₂/kg/hr, respectively. Seasonal metabolic compensation was compared in spotted seatrout and sand seatrout in Redfish Bay, Texas (Vetter 1982). Spotted seatrout controlled their metabolic rates within a narrower range in response to seasonal temperature change (50 mg O₂/kg/hr at 15°C, 124 mg O₂/kg/hr at 30°C) than did sand seatrout (25 mg O₂/kg/hr at 15°C, 170 mg O₂/kg/hr at 30°C). Sand seatrout migrate from the estuaries to the Gulf of Mexico in late summer and overwinter there, whereas spotted seatrout are permanent residents of estuaries. Vetter (1982) suggested that greater metabolic compensation on the part of spotted seatrout may be an adaptation to year-round exploitation of the estuarine habitat, which has more extreme temperatures than offshore waters.

12.3 Life History

Various aspects of spotted seatrout life history were reviewed by Guest and Gunter (1958), Futch (1970), Idyll and Fahy (1970), Lorio and Perret (1980), and Perret et al. (1980).

12.3.1 Reproduction

Spotted seatrout mature between one and three years of age and males tend to mature at a smaller size than females (Table 12-2). Size at maturity varies from estuary to estuary (Lorio and Perret 1980; Perret et al. 1980).

Estimates of spotted seatrout fecundity ranged from 1.4×10^4 to 1.6×10^7 (Table 12-3). Sundararaj and Suttkus (1962) concluded that age III fish had the greatest "spawning power", producing 40.6% of the egg supply followed by age IV (26.8%) and age II (24.5%) fish. Overstreet (1983) treated all oocytes >30 μ m (other cited authors counted only large yolky eggs) because spawning occurs over several months and because vitellogenesis can proceed rapidly in small oocytes.

Table 12-2. Reported size and/or age at maturity for spotted seatrout. Number in parentheses is the smallest reported size or age at maturity.

Reference	Locality	Size			Age	
		Males	Females	Both	Males	Females
Pearson (1929)	Texas					2-3
Miles (1950)	Texas			200-250mm SL		
Hein and Shepard (1979a)	Louisiana	(163mm SL)	(207mm SL)			
Overstreet (1983)	Mississippi		220-299mm SL (140-219mm SL)			
Klima and Tabb (1959)	Northwest Fla.	25cm SL (18cm SL)	27cm SL			
Moody (1950)	Cedar Key, Fla.	200-240mm SL	210-250mm SL			
Stewart (1961)	Southwest Fla.			290-300mm SL (190mm SL)		
Rutherford et al. (1982)	Southwest Fla.	(237mm SL)	(230mm SL)		2-4	3-4
Tabb (1961)	East Central Fla.	35cm SL	35cm SL		2-3	3-4
Music and Pafford (1984)	Georgia	(342mm TL)	(229mm TL)			(2)
Brown (1981)	Chesapeake Bay	250mm TL	290-350mm TL		2	3

Table 12-3. Fecundity estimates for spotted seatrout as reported in the literature. [*Total length approximated by TL=10.6586+1.1284SL from Overstreet (1983)]

Reference	Locality	Average Size (mm TL)	N	Calculated Number of Eggs	Egg Diameter (mm)	Method
Pearson (1929)	Texas	480	1	427,819	not given	gravimetric
		620	1	1,118,000		
Moody (1950)	Cedar Key, Florida	459*	1	464,000	not given	gravimetric
Tabb (1961)	Indian River area, Florida	377*	3	15,000	not given	not given
		509*	3	150,000	given	
		575*	3	400,000		
		716*	3	1,100,000		
Sundararaj and Suttus (1962)	Louisiana	283	8	140,485	.38	gravimetric (mean egg diameter)
		376	9	354,325	.52	
		450	8	660,960	.61	
		504	3	1,144,492	.63	
Overstreet (1983)	Mississippi Sound	379*	1	2,254,134	.30	volumetric
		595*	1	3,968,050	.30	gravimetric
				10,599,376	.03	volumetric
				15,567,833	.03	gravimetric

The spotted seatrout has a protracted spring and summer spawning season which peaks in late April-July in the Gulf of Mexico (Lorio and Perret 1980; Perret et al. 1980). Spotted seatrout spawn from April through September along the Atlantic coast. Spawning off east-central Florida occurs from mid-April to September (Tabb 1961; Gilmore et al. 1976; Mok and Gilmore 1983). In Georgia spotted seatrout spawn from April to August with a peak in May (Mahood 1975) and second smaller peak in July (Music and Pafford 1984). Limited collections of larvae along the Carolina coasts indicated an April-August spawning season (Hildebrand and Cable 1934; Powles and Stender 1978). Two spawning peaks were observed in Chesapeake Bay, mid-May to mid-June and July, corresponding to early maturing and late maturing groups of fish (Brown 1981).

Brown et al. (1983) compared reproductive strategies for spotted seatrout in Redfish Bay, Texas and Chesapeake Bay, Virginia. Spotted seatrout are year-round residents of Redfish Bay and spawn for seven months (late March-early April through October), whereas they only reside in Chesapeake Bay from May through October and spawn from May to August. In Texas all sexually mature males and 80% of mature females were ripe throughout the spawning season. In Virginia only 25% of the mature males were ripe throughout the spawning season and ripe females were only found during spawning peaks. Histological observations suggest multiple spawning in Texas. Maturity is reached one year earlier in Texas than in Virginia.

The preferred spawning habitat of spotted seatrout has not been precisely determined but may include offshore and estuarine areas. Spawning in the Gulf of Mexico is believed to occur in the deeper portions (3.0 - 4.6 m) of bays and lagoons over grassy areas (Pearson 1929; Miles 1950; Moody 1950; Stewart 1961; Tabb and Manning 1961; Tabb 1966; Overstreet 1983) and in the inshore waters of the Gulf along barrier islands, particularly in or near coastal passes (King 1971; Jannke 1971; Christmas and Waller 1973; Sabins and Truesdale 1974; Hein and Shepard 1979a; Houde et al. 1979; Overstreet 1983).

Spawning on the Atlantic coast probably occurs in coastal and estuarine waters. Tabb (1961) believed that spawning in the Indian River lagoon system, Florida took place in the deeper channels immediately adjacent to the vegetated shallows with a dispersion of young to the shallow grassy bays with increasing size. This is supported by Mok and Gilmore's (1983) study of spotted seatrout sound production in the Indian River lagoon, Florida, which found that the highest intensity of large group sounds was limited to the Intracoastal Waterway and adjacent deeper parts of the seagrass flats. Small group and individual sounds appeared on both sides of the Intracoastal Waterway and extended into shallow seagrass areas. Acoustic activity and egg numbers (indicating spawning) were positively correlated in that study. Spotted seatrout in Georgia spawn along beaches near tidal inlets and mouths of sounds, and within creeks and sounds, usually in water 0.9-3.0 m deep (Mahood 1975; Music and Pafford 1984). Spawning along the South Carolina coast probably takes place in the lower portions of estuaries and inlets (Powles and Stender 1978). Hildebrand and Cable (1934) were unable to determine the exact spawning ground(s) of spotted seatrout in North Carolina based on limited collections of larvae taken from 8-10 km

offshore to within an estuary. Running ripe fish were caught over seagrass beds in Chesapeake Bay within several hundred feet of a channel (Brown 1981).

Spotted seatrout spawn at night (Tabb 1966; Brown 1981; Mok and Gilmore 1983). Holt et al. (1983) reported synchronous spawning near dusk in Texas. During spawning there is a constant milling and jumping of the spawning school, with side-to-side body contact among the fish (Miles 1950; Tabb 1966). Miles (1950) examined several spent males and found that they were rubbed raw around the pelvic fins, lower abdomen and vent. Spawning is accompanied by croaking sounds produced only by the males (Smith 1907; Stewart 1961; Tabb 1966). The drumming muscles of mature males (>193 mm TL) were deeper red during the spawning season than at other times of the year (Hein and Shepard 1979a). Croaking was generally heard approximately one to two hours before sunset and continued for up to six hours (Mok and Gilmore 1983). Sound production has been noted at times other than spawning, suggesting a secondary function such as defense (Stewart 1961; Hein and Shepard 1979a).

Spotted seatrout spawn at temperatures from 21-28°C. Spawning was reported to occur at 21°C or higher in Texas (Simmons 1951) and from 21-35°C in Louisiana (Fontenot and Rogillio 1970; Rogillio 1975; Hein and Shepard 1979a). Stewart (1961) stated that bimodal peaks of capture of ripe adults corresponded with the 28-30°C temperature range in southwestern Florida. Jannke (1971), however, stated that a temperature of 24°C or greater appears necessary to initiate spring spawning in southwestern Florida. Tabb (1966) reported that spawning took place between 25.5 and 28.3°C on the east-central coast of Florida.

Hein and Shepard (1979a) reported that peak spawning in Louisiana occurred in May in 1976 and 1978 on an increasing photoperiod, while the second major peaks (July 1976 and August 1978) were recorded on a decreasing photoperiod for both years. The amount of daylight during which gravid fish were collected remained nearly the same for both years: 13 hr, 42-59 min, and 13 hr, 10-41 min (sunrise to sunset).

Fluctuating salinity is a common factor of all productive spotted seatrout grounds (Tabb 1966). Peak spawning in Florida waters occurred when salinities reached 30-35 ppt in the lagoons and estuaries during dry spring months. Hein and Shepard (1979a) collected gravid spotted seatrout at salinities of 21-26 ppt in 1976 and 17-26 ppt in 1978. No spawning occurred in the Laguna Madre of Texas when salinity exceeded 45 ppt (Simmons 1957).

Arnold et al. (1976) found that spotted seatrout spawn in the laboratory over a salinity range of 25-30 ppt at 26°C, coupled with a constant daylight period of 15 hours.

Laboratory-spawned spotted seatrout eggs were pelagic and spherical with an average diameter of 0.77 mm and usually contained one yellow oil globule (2% had two to three globules) (Fable et al. 1978). Miles (1950, 1951) reported that eggs from ripe ovaries measured 0.70-0.98 mm in diameter and contained one to four small oil globules. Tabb (1966) stated that eggs were spherical and normally had one oil droplet, but sometimes two or three.

12.3.2 Pre-adult Phase

Fable et al. (1978) described embryonic development of reared eggs (Figure 12-2). Hatching occurred 16-20 hr after fertilization at incubation temperatures of approximately 25°C. Smith (1907) reported that spotted seatrout eggs hatched in 40 hr at 25°C.

Larval and juvenile development of spotted seatrout was described and illustrated by Welsh and Breder (1923), Pearson (1929), Hildebrand and Cable (1934), and Jannke (1971) (Figure 12-2). These studies were recapitulated by Lippson and Moran (1974) and Johnson (1978). Daniels (1977) described and illustrated larvae 1.8-11.3 mm SL. Fable et al. (1978) described the larval development and morphometrics of laboratory-reared spotted seatrout from hatching (1.3-1.5 mm SL) to 15 days (4.5 mm SL). Descriptions and morphometrics of 25 specimens (1.9 - 32.2 mm SL) from South Carolina were presented and compared with previous studies by Powles and Stender (1978).

12.3.3 Adult Phase

Age and growth studies of spotted seatrout indicate that longevity is greater in the northern part of the range. Brown (1981) found age XV (776 mm TL) to be the maximum age in a Chesapeake Bay study. The maximum age for east-central Florida spotted seatrout was age X (Tabb 1961). Age VIII spotted seatrout were reported from Georgia (Music and Pafford 1984) and the Gulf coast of Florida (Moffett 1961). Pearson (1929) reported several age IX fish in Texas. The largest spotted seatrout reported in the literature were two fish weighing 7.25 kg caught in Chesapeake Bay in 1922 and at Mason's Beach, Virginia in 1977 (Hildebrand and Schroeder 1928; Anonymous 1983a), and a 7.48 kg fish caught in the Neuse River in 1903 (Smith 1907). Using Brown's (1981) length-weight relationship these individuals would measure 875 and 884 mm TL, respectively.

Nine predators and six competitors of spotted seatrout were listed for Apalachee Bay in northwest Florida (Table 12-4), whereas only five predators, spotted seatrout, snook, gafftopsail catfish, common jack, and mangrove snapper, were listed for east-central Florida (Klima and Tabb 1959). Tabb (1961) suggested that the scarcity of predator and competitor species was a factor in the abundance and "success" of spotted seatrout in estuarine habitats.

Parasites, diseases, mortalities, and abnormal conditions of spotted seatrout were reviewed by Perret et al. (1980) and Overstreet (1983).

12.3.4 Nutrition and Growth

Spotted seatrout are carnivorous, feeding primarily on crustaceans and fishes (Linton 1905; Hildebrand and Schroeder 1928; Pearson 1929; Gunter 1945; Kemp 1949; Knapp 1949; Miles 1949; Moody 1950; Reid 1954; Darnell 1958; Klima and Tabb 1959; Springer and Woodburn 1960; Stewart 1961; Tabb 1961; Lorio and Schafer 1966; Seagle 1969; Fontenot and Rogillio 1970; Odum 1971; Adams 1972; Carr and Adams 1973; Mahood 1975; Odum and Heald 1972; Rogillio 1975; Orth and Heck 1980; Weinstein 1981;




Figure 12-2. Stages in the development of spotted seatrout embryos, larvae, and juveniles (illustration from Lippson and Moran 1974; Fable et al. 1978).

Table 12-4. List of predators and competitors of spotted seatrout in Apalachee Bay in northwest Florida (from Klima and Tabb 1959).

Common name	Scientific name	Relation	Occurrence
Striped bass	<u>Roccus saxatilis</u> (Walbaum)	Predator (?)	Resident
Snook	<u>Centropomus undecimalis</u> (Bloch)	Predator	Seasonal
Tarpon	<u>Megalops atlantica</u> Valenciennes	Predator	Seasonal
Alligator gar	<u>Lepisosteus spatula</u> Lacepede	Predator (?)	Seasonal
Sea catfish	<u>Galeichthys felis</u> (Linnaeus)	Competitor	Resident
Barracuda	<u>Sphyraena barracuda</u> (Walbaum)	Predator	Occasional
Spanish mackerel	<u>Scomberomorus maculatus</u> (Mitchill)	Predator	Seasonal
King mackerel	<u>Scomberomorus cavalla</u> (Cuvier)	Predator	Seasonal
Bluefish	<u>Pomotomus saltatrix</u> (Linnaeus)	Predator	Resident
Grouper	<u>Mycteroperca</u> sp.	Competitor	Resident
Silver perch	<u>Bairdiella chrysura</u> (Lacepede)	Predator and competitor	Resident
Red drum	<u>Sciaenops ocellatus</u> (Linnaeus)	Competitor	Resident
Spot	<u>Leiostomus xanthurus</u> (Lacepede)	Competitor	Resident
Croaker	<u>Micropogon undulatus</u> (Linnaeus)	Competitor	Resident
Southern rock bass	<u>Ambloplites rupestris</u> <u>arionus</u> Viosca	Competitor	Resident

Overstreet and Heard 1982; Rutherford et al. 1982; Matlock and Garcia 1983). The most important crustaceans were penaeid shrimp and crabs. Anchovies, menhaden, mullet, pinfish, and silversides accounted for the highest percentage of fishes in spotted seatrout stomach contents (Table 12-5).

Changes in food habits with growth were noted in several studies (Moody 1950; Darnell 1958; Tabb 1961; Adams 1972; Carr and Adams 1973; Colura et al. 1976). Copepods were important in the diet of fish <30 mm TL, and mysids, caridean shrimp, palaemonid shrimp, amphipods, polychaetes, and aquatic insects were important in fish <150 mm. Crustaceans (penaeid shrimp and blue crabs) were more important in fish 150-275 mm SL, whereas fish (pinfish, mullet, anchovies, and menhaden) predominated in larger fish (Moody 1950; Seagle 1969; Overstreet and Heard 1982).

Tabb (1961) suggested that food preferences are probably the result of seasonal availability of food. In Florida waters shrimp are most abundant during summer and early winter and fishes are more abundant in late winter and early spring. Lorio and Schafer (1966) also noted that shrimp were most available and eaten more frequently during summer by spotted seatrout in Louisiana. The food contents of fish obtained from Mississippi Sound included a slightly greater percentage of fish during spring and summer, when anchovies were more common. Penaeids were less prevalent in stomach contents during autumn and winter when they were also less available in the study area (Overstreet and Heard 1982).

A high percentage of empty stomachs were reported in most of these studies (Table 12-5). Moody (1950) attributed this to sporadic feeding. Darnell (1958) and Seagle (1969) suggested that spotted seatrout feeding was heaviest during the early to mid-morning hours. Lascara (1981) reported movements of spotted seatrout in and out of submerged grass beds in Chesapeake Bay and found times of peak abundance corresponded to crepuscular periods (dawn and dusk). He believed that these were times of maximum feeding. Spotted seatrout have a tapeta lucida in their eyes which allows greater light penetration to the retina (Arnott et al. 1971). This adaptation allows for keener eyesight than potential prey items in dim light, and may increase feeding duration and success during periods of low light intensity. Experienced, highly successful spotted seatrout fishermen in Virginia reported that best fishing is at dusk, drifting over grass beds on a flood tide (Brown 1981).

Growth of spotted seatrout larvae in the laboratory increased significantly as prey concentrations and temperatures were raised, and decreased as stock density increased (Taniguchi 1979, 1981; Houde and Taniguchi 1982). Juvenile spotted seatrout (100-112 mm TL), stocked in ponds in Louisiana, grew 2.08 mm/day in October and 0.33 mm/day in November (Sackett et al. 1979).

Spotted seatrout growth is rapid during the first year. Pearson (1929) reported a modal length of 130 mm TL in Texas by the first winter with a range of 50-200 mm TL which reflects the prolonged spawning period. In Georgia juvenile spotted seatrout attained a mean length of 124 mm TL in November (Mahood 1975). Hildebrand and Cable (1934) reported a modal length of 170 mm TL for juvenile spotted seatrout in North Carolina at

the end of 7-8 months of growth. Welsh and Breder (1923) collected five juveniles from Chesapeake Bay in December ranging in length from 110-125 mm TL. Brown (1981) reported that spotted seatrout in Chesapeake Bay attained an average of 170 mm TL by the end of their first winter. Spotted seatrout growth slows considerably by age II. Differences in age and growth between the sexes and between different populations are discussed in Section 12.4.1.

12.3.5 Behavior

Tagging studies indicated that spotted seatrout are relatively non-migratory in Georgia, Florida, and the Gulf of Mexico. Although no tagging studies have been done north of Georgia, indications are that spotted seatrout migrate seaward and southward from Chesapeake Bay and possibly from North Carolina sounds (Section 12.2.2.2). Most movement in Georgia estuaries was short range, averaging 8.9 km, although two individuals traveled 105 and 110 km, respectively (Music 1981; Music and Pafford 1984). Movement is apparently restricted to seasonal migrations in and out of the open sounds enroute to creeks and rivers in fall and winter, and off the beaches in spring and early summer. Moffett (1961) reported that over 95% of all returned tagged fish on the west coast of Florida were recovered within 48 km of the tagging sites. The longest migration was 507 km, from Apalachicola, Florida to Grand Isle, Louisiana. Results of other tagging studies in Florida also indicated that spotted seatrout rarely move over 32-48 km (Ingle et al. 1962; Topp 1963; Beaumariage 1964, 1969; Beaumariage and Wittich 1966).

Tabb (1966) reported that spotted seatrout begin to school by the age of 6 to 8 weeks (25-50 mm). Schooling behavior remains pronounced until an age of about 5 to 6 years, at which time most males have died and the remaining large females (2.7-3.6 kg), called "sow" or "gator" trout, adopt a semi-solitary existence.

12.3.6 Contaminants

Trace element levels were determined for 15 elements in spotted seatrout to provide baseline data to help identify potential problems involving species, elements, or locations (Hall et al. 1978). No interpretive comments were provided.

The acute lethal effects of sodium hypochlorite, chloramine, and 5-chlorouracil on eggs and larvae of spotted seatrout were presented by Johnson et al. (1977). Forty-eight-hour median tolerance limits (TLM) for the various toxicants and age classes (2-hr old eggs, 10-hr eggs, and 1-hr posthatch larvae) were, respectively: sodium hypochlorite - 0.21 ± 0.01 , 0.21 ± 0.01 , 0.17 ± 0.28 ppm; chloramine - 14.14 ± 1.13 , 0.57 ± 0.28 , 5.75 ± 3.01 ppm; and 5-chlorouracil - 8.91 ± 1.03 , >100 , 79.43 ± 44.97 ppm. Results of this study indicated that considerable larval seatrout loss would be expected in areas of chlorinated effluent disposal where the toxic products of sodium hypochlorite and seawater are 0.17 ppm sodium hypochlorite.

Sublethal effects of fuel oil (water-soluble fraction) on larval spotted seatrout were investigated by Johnson et al. (1979). When larvae were

Table 12-5. Stomach contents of spotted seatrout from different estuarine areas along the U.S. Atlantic and Gulf of Mexico coasts.

Author	Mahood (1975)	Tabb (1961)	Rutherford et al. (1982)
Locality	Georgia	Indian River, Florida	Everglades National Park
Period	Oct. 1970-Sept. 1973	Oct. 1955-June 1957	Nov. 1978-June 1980
Source	Table 6, p. 206	Table 6, p. 20	Figure 16, p. 38
Number of specimens	108	580	724
Empty stomachs		410	486
Length of specimens		2.5-15 cm SL	15-85 cm SL
Quantitative method	% of occurrence	% total number of food items	< 271 mm 271-370 mm > 370 mm
			% of occurrence
Fishes:			
Engraulidae	5.9	-	¹ 20.0 ¹ 46.0 ¹ 58.3
Clupeidae	10.2	-	-
Mugilidae	5.9	-	1.3
Atherinidae	7.6	-	22.5
Cyprinodontidae	5.1	-	2.6
Sparidae	-	-	10.6
Sciaenidae	8.3	-	3.3
Other	2.4	-	3.3
Unidentified fish	8.5	3.6	4.6
Crustaceans:			
Mysidacea	-	42.8	-
Amphipoda	-	-	-
Isopoda	-	-	-
Penaeid shrimp	33.9	53.6	88.3 72.0 58.3
Caridean shrimp	2.5	-	-
Brachyuran crabs	-	-	1.0 4.0 8.0
Other invertebrates:	9.3	-	-
Polychaetes	-	-	-
Mollusks	-	-	5.0 12.0 -
Vegetation			12.0 20.0 5.0
Unidentified remains	-	-	2.0 5.0 5.0
¹ All fishes combined			

Table 12-5. (continued)

Author	Overstreet and Heard (1982)	Lorio and Schafer (1966)	Seagle (1969)
Locality	Mississippi Sound	Biloxi Marsh Area, La.	Redfish Bay, Texas
Period		1961 - 1965	1967 - 1968
Source	Table 2, p. 143	Table II, p. 292	Table 1
Number of specimens	111	368	205
Empty stomachs	11	152	39
Length of specimens	73-249	25-51 cm TL	226-350
Quantitative method	% of occurrence	% of occurrence	% of occurrence
Fishes:	¹ 65.0	¹ 74.4	
Engraulidae		1.4	20.0
Clupeidae		1.4	25.4
Mugilidae		0.4	8.0
Atherinidae		-	2.3
Cyprinodontidae		-	4.0
Sparidae		-	-
Sciaenidae		0.9	14.0
Other		0.9	4.0
Unidentified		68.5	2.0
Crustaceans:	² 47.0	² 25.3	
Mysidacea		-	1.5
Amphipoda		0.9	3.9
Isopoda		2.3	1.5
Penaeid shrimp		11.1	40.0
Caridean shrimp		6.5	23.1
Brachyuran crabs		4.1	1.5
Other invertebrates:			6.2
Polychaetes	2.0	9.8	8.0
Mollusks	2.0	1.3	-
Vegetation	3.0	6.7	-
Unidentified remains	-	-	-
¹ All fishes combined			
² All crustaceans combined			

subjected to sublethal concentrations of fuel oil (0.00-1.00 ppm), there was a general decrease in total body length and critical distance, while the percentage of larvae with unpigmented eyes increased with increased oil concentration.

DDT residues were measured in six generations of spotted seatrout from the Laguna Madre, Texas (Butler 1969; Butler et al. 1970). Residues reached as high as 8 ppm in the gonads and breeding apparently did not occur for at least one or two years.

A survey of polychlorinated biphenyls (PCBs) in selected finfish species determined that the mean level of PCBs in 17 spotted seatrout from the Gulf of Mexico was 0.16 ppm. This level is far below existing (5 ppm) or proposed (2 ppm) maximum permissible levels in foodfish (Gadbois and Maney 1983).

12.4 Population

12.4.1 Structure

Aspects of spotted seatrout population structure have been investigated for Virginia (Brown 1981), Georgia (Mahood 1974, 1975; Music and Pafford 1984), Florida (Tabb 1958, 1961; Klima and Tabb 1959; Moffett 1961; Stewart 1961; Iversen and Tabb 1962; Rutherford 1982; Rutherford et al. 1982), Alabama (Tatum 1980; Wade in press), Louisiana (Hein and Shepard 1979a, 1980), and Texas (Pearson 1929).

Reported sex ratios for spotted seatrout populations indicate that overall, females outnumber males by as much as three to one. Pearson (1929) reported a 2:1 ratio favoring females in Texas. Female spotted seatrout outnumbered males 2.4:1 in a Louisiana study (Hein and Shepard 1979a). In western Florida females dominated at all ages and ratios increased with age to as much as 9:1 at age V (Klima and Tabb 1959; Moffett 1961). Overall sex ratios were 2:1 for both east and west coast Florida populations (Tabb 1961). Rutherford et al. (1982) found an overall sex ratio favoring females by 1.7:1, which remained constant at ages II through V in Everglades National Park, Florida. In Georgia studies the overall sex ratio of females to males were 1.7:1 (Mahood 1975) and 1.9:1 (Music and Pafford 1984). The ratio changed from 1:3 for fish <250 mm to 1:1 for fish 251-350 mm, 2.6:1 for fish 351-400 mm, and 23:1 for fish 501-550 mm TL (Music and Pafford 1984).

Older spotted seatrout were found in Chesapeake Bay (age XV) than from populations to the south (Brown 1981). Tabb (1961) reported age X fish in east-central Florida and Pearson (1929) found age IX spotted seatrout in Texas. Age VIII was the maximum reported age for Georgia (Music and Pafford 1984) and Fort Meyers, Florida (Moffett 1961). Age VII spotted seatrout were reported for southwestern Florida (Stewart 1961; Rutherford 1982) and northwest Florida (Klima and Tabb 1959). Moffett (1961) found fish through age VI in west Florida (Cedar Key), as did Tatum (1980) and Wade (in press) in Alabama.

The age distribution of the catch varied in different areas. Age groups III (27%) and IV (21%) dominated the catch in Chesapeake Bay (Brown 1981). These samples were mainly collected by haul seines which equally sample all sizes of the recruited population. Tabb (1961) found that age groups I (28%), II (27%) and III (21%) predominated in his samples from east-central Florida. In southwestern Florida dominant ages apparently shifted from age II (36%) and age III fish (42%) in 1959 (Stewart 1961) to age III (45%) and age IV (29%) in 1979 (Rutherford 1982). In west Florida age groups II (41%) and III (35%) predominated (Moffett 1961). Age group III (47%), followed by ages II (23%) and IV (22%), predominated in a northwest Florida study (Klima and Tabb 1959). Difference between the Florida studies were caused in part by differences in gear selectivity, while the lack of younger fish in these studies reflected a minimum legal size limit (30.5 cm). Only Tabb (1961) and Stewart (1961) obtained smaller spotted seatrout by trawling. Tatum (1980) reported that age groups III+ (27.7%) and II+ (17.8%) were the two most exploited age classes in Alabama fishing tournaments, 1964-1977; however, age groups V+, VI+ and >VI+ were most abundant in 1965, 1970, and 1971. Age III spotted seatrout, followed by II and IV, dominated the catch in 3-1/4 in stretched mesh gill nets in a Louisiana study (Hein and Shepard 1980).

Female spotted seatrout attain a greater maximum age than males throughout the range (Table 12-6). Brown (1981) found that males and females reached at least ages VIII and XII, respectively, in Chesapeake Bay. The sex of the age XV fish collected in that study was not determined. The oldest male and female spotted seatrout aged in Georgia were ages VI and VIII, respectively (Music and Pafford 1984). All Florida studies found that females lived at least one year longer than males (Klima and Tabb 1959; Moffett 1961; Stewart 1961; Tabb 1961; Rutherford 1982) except at Cedar Key, Florida (Moffett 1961). The tendency of female spotted seatrout to outlive males was also reported for Alabama (Wade in press), Louisiana (Hein and Shepard 1979a) and Texas (Pearson 1929).

Age and growth studies of spotted seatrout revealed that size at age varies between locations (Table 12-6). Reported mean back-calculated lengths at age were largest for east-central Florida (Tabb 1961). There appears to be a discrepancy in Tabb's data, since his back-calculated lengths at age for combined sexes were considerably greater than lengths for either males or females, separately (Table 12-6). Spotted seatrout aged in Georgia were larger (combined sexes) than Chesapeake Bay fish at each age except age VI, and smaller than east-central Florida fish (Music and Pafford 1984). Calculated lengths for ages I and II spotted seatrout in Everglades National Park were greater than reported in other studies because of the back calculation formula used: $L_t - a = S_t (L - a) / S$, where L_t = length at annulus t , L = length of fish at capture, S_t = scale radius at annulus t , S = total scale radius, and a = y intercept of fish length regressed on total scale radius (Rutherford 1982). Previous investigators calculated fish lengths at annulus directly according to the formula: $L_t = S_t L / S$. The difference in length resulting from the type of back-calculation formula used becomes negligible after age II when lengths of the Park spotted seatrout population closely parallel lengths reported for other

Table 12-6. Mean back-calculated standard lengths (mm) at age for spotted seatrout as reported in the literature.

Study	Location	Sex	I	II	III	IV	Age					XI	XII
							V	VI	VII	VIII	IX		
Brown (1981)*	Chesapeake Bay, Va.	Male	128 (68)	206 (66)	267 (58)	323 (26)	370 (20)	430 (7)	430 (7)	475 (2)			
		Female	161 (92)	237 (91)	296 (80)	346 (57)	385 (35)	449 (16)	449 (16)	481 (12)	504 (4)	553 (1)	
		Combined	143 (247)	221 (221)	302 (193)	354 (132)	378 (60)	447 (35)	447 (35)	482 (21)	518 (6)	557 (5)	620 (2)
Music and Pafford (1984)*	Georgia	Male	148 (213)	228 (219)	280 (109)	323 (26)	363 (13)	385 (9)					
		Female	161 (333)	256 (326)	328 (227)	386 (108)	425 (57)	463 (27)	523 (7)	572 (2)			
		Combined	155 (548)	244 (538)	313 (336)	373 (134)	413 (70)	443 (36)	523 (7)	572 (2)			
Tabb (1961)	East-Central, Fla.	Male	138 (281)	200 (195)	259 (109)	322 (43)	380 (14)	410 (4)	437 (1)				
		Female	145 (682)	220 (509)	300 (336)	353 (183)	408 (76)	480 (27)	520 (16)	545 (2)			
		Combined	165 (592)	248 (563)	317 (436)	384 (234)	457 (135)	533 (73)	561 (36)	624 (1)			
Stewart (1961)	Flamingo, Fla.	Male	128 (433)	210 (417)	267 (217)	313 (33)	348 (5)						
		Female	137 (562)	236 (538)	280 (372)	345 (133)	401 (50)	434 (10)	451 (2)				
Rutherford (1982)	Everglades National Park, Fla.	Male	215 (205)	260 (202)	292 (164)	327 (65)	377 (12)	403 (2)					
		Female	210 (326)	264 (325)	307 (275)	354 (134)	402 (32)	420 (12)	488 (3)				
		Combined	212 (539)	262 (535)	301 (446)	345 (201)	395 (44)	417 (12)	488 (3)				
Moffett (1961)	Fort Meyers, Fla.	Male	128 (258)	206 (235)	259 (154)	310 (39)	354 (13)	437 (1)					
		Female	131 (386)	209 (383)	266 (293)	322 (118)	371 (46)	409 (21)	431 (5)	438 (1)			
		Combined	130 (644)	208 (638)	264 (447)	320 (157)	368 (59)	430 (22)	431 (5)	438 (1)			
Moffett (1961)	Cedar Key, Fla.	Male	129 (83)	206 (83)	263 (52)	323 (19)	380 (4)	434 (1)					
		Female	130 (217)	212 (215)	269 (151)	323 (48)	383 (5)						
		Combined	130 (300)	211 (298)	268 (203)	323 (67)	382 (9)						
Klima and Tabb (1959)	Apalachicola, Fla.	Male	115 (316)	188 (312)	250 (261)	304 (71)	341 (7)	369 (1)					
		Female	117 (549)	191 (549)	258 (525)	315 (251)	372 (66)	423 (8)	437 (1)				
		Combined	116 (665)	190 (661)	255 (786)	312 (322)	369 (73)	422 (9)	437 (1)				
Pearson (1929)*	Texas	Combined	123 (396)	203 (303)	259 (230)	305 (160)	340 (94)	377 (57)	418 (36)	444 (8)	488 (3)		

*TL converted to SL using $SL = -3.8832 + 0.8653TL$ (Hain et al. 1980)

populations (Rutherford 1982). Spotted seatrout growth in Chesapeake Bay appears similar to growth at Flamingo and Everglades National Park (excluding ages I and II). Brown (1981) reported that growth in Chesapeake Bay for the population as a whole was significantly different from growth in Texas and east-central Florida, but not from Fort Meyers, Cedar Key, or Apalachicola, Florida.

Female spotted seatrout were larger than males for all ages at each location except for age I fish at Everglades National Park and age VI fish at Fort Meyers, Florida where there was a single large age VI male (Table 12-6). Growth of males and females in Chesapeake Bay was significantly different from growth of Fort Meyers, Cedar Key, and Apalachicola, Florida fish, although growth for the population as a whole was not significantly different (Brown 1981).

Von Bertalanffy growth parameters were derived for spotted seatrout in Chesapeake Bay (Brown 1981) and Everglades National Park (Rutherford 1982) (Table 12-7). Brown's (1981) values of t_0 and K appear too low and the resulting growth curves do not describe growth as determined by back-calculated lengths. The L_∞ values for females are higher than for males as would be expected, since females attain a greater length at age than males. Rutherford's (1982) growth parameters derived by Bayley's (1977) method, appear to closely predict spotted seatrout lengths at ages I-VI. The lower L_∞ values for Everglades National Park fish reflect the fewer age classes and smaller sizes present in the samples.

Length-weight relationships were determined for spotted seatrout in Texas (Harrington et al. 1979), Louisiana (Adkins et al. 1979; Hein et al. 1980), Mississippi (Overstreet 1983), Alabama (Wade in press), Florida (Moffett 1961; Rutherford 1982), and Virginia (Brown 1981) (Table 12-8). Spotted seatrout from Chesapeake Bay appear to be heavier at a given length than those from other areas. Brown (1981), however, stated that these fish were collected only during the summer at the period of maximum feeding and sexual activity when weights would tend to be higher. No significant differences in length-weight relationships between sexes were found in Alabama or Everglades National Park, although males appeared to be heavier than females at all lengths in Alabama (Wade in press; Rutherford 1982).

Age and growth differences, the non-migratory nature of spotted seatrout and the isolation of estuarine areas along the South Atlantic and Gulf coasts suggest that there are distinct subpopulations of spotted seatrout. Iversen and Tabb (1962) concluded that there were separate populations in Florida based on growth and tagging data. Electrophoretic studies by Weinstein and Yerger (1976b) supported the concept of genetically distinct populations of spotted seatrout in the estuaries they sampled. The populations sampled west of the Mississippi River formed a group distinct from those populations east of the Mississippi. Within each of these regions were separate populations, such as Corpus Christi and Galveston Bay to the west, and St. Joseph, Apalachee, Tampa, and Florida Bays in the east. Florida Bay was the most divergent of eastern Gulf populations, possibly because of the unique environmental characteristics of this estuary (shallow mud flats with higher turbidities and higher average yearly temperatures). The

Table 12-7. Von Bertalanffy growth parameters as reported in the literature.

Area	Sex	L_{∞} (mm)	t_0	K
Chesapeake Bay, Va.	Male	760	-7.5933	.052
	Female	854	-7.9278	.052
	Combined	935	-5.6091	.059
Everglades National Park, Florida	Male	591	-2.95	.12
	Female	656	-2.04	.13
	Combined	774	-2.54	.09

¹Brown 1981²Rutherford 1982

Table 12-8. Length-weight relationships for spotted seatrout collected from different localities.

Locality	Sex	N	Length range (mm-TL)	Equation	r	Calculated weight for a 350 mm TL fish
Chesapeake Bay, Virginia ¹	Combined	340	122-782	$\log W = -5.072 + 3.043 \log TL$	0.99	469
	Males	74	264-580	$\log W = -5.598 + 3.244 \log TL$	0.99	448
	Females	102	135-760	$\log W = -4.924 + 2.986 \log TL$	0.97	470
Georgia ²	Combined	710		$\log W = -4.848 + 2.949 \log L$	0.95	451
	Males	277		$\log W = -4.182 + 2.683 \log L$	0.86	440
	Females	409		$\log W = -4.516 + 2.824 \log L$	0.93	466
Everglades National Park, Florida ³	Combined	567		$\log W = -5.194 + 2.745 \log SL$	0.93	400
Fort Myers and Cedar Key, Florida ⁴	Combined	307		$\log W = -5.333 + 3.113 \log TL$		386
Alabama ⁵	Combined			$\log W = -5.305 + 3.105 \log TL$		392
Mississippi ⁶	Males	1,087	40-465(SL)	$\log W = -4.947 + 3.051 \log SL$	0.99	404
	Females	1,683	39-544(SL)	$\log W = -4.947 + 3.052 \log SL$	0.99	406
South-Central, Louisiana ⁷	Combined	1,208	21-629	$\log W = -5.423 + 3.154 \log TL$	0.99	400
Texas ⁸	Combined	9,498	49-902	$\log W = -5.192 + 3.062 \log TL$	0.99	396

¹ Brown 1981² Music and Pafford 1984³ Rutherford 1982⁴ Moffett 1961⁵ Wade (In press)⁶ Overstreet 1983⁷ Hein et al. 1980⁸ Harrington et al. 1979

most widely divergent groups were those populations west of the Mississippi River and on the Atlantic coast of Florida (Indian River).

12.4.2 Abundance, Density, Mortality, and Dynamics

Peaks in spotted seatrout abundance occur in spring and/or fall and winter in various estuaries throughout the range. Adults are particularly abundant in Texas and Florida in spring when they migrate from overwintering areas through passes and channels to shallow feeding and possibly spawning areas (Pearson 1929; Tabb 1958). In the lower Laguna Madre, Texas, adults averaged 2.58 kg/ha in spring in 1970-72, in contrast to 1.52 kg/ha during fall (Breuer 1973). Adult spotted seatrout are also relatively abundant during winter when they concentrate in deeper holes to escape cold (Pearson 1929; Gunter 1938; Perret 1971; Waller and Sutter 1982). Highest catch per unit effort of spotted seatrout in the shallow (0.6-1.2 m) Biloxi Marsh complex of Louisiana occurred in fall with a smaller peak in spring (Fontenot and Rogillio 1970). Adkins et al. (1979) reported that peak abundance in Louisiana occurred in spring with a second peak in late summer or fall. In Georgia spotted seatrout were most abundant in the deeper waters of the sounds and creeks in winter and in the shallows in spring (Mahood 1975). Brown (1981) reported that recreational catches of spotted seatrout in Chesapeake Bay were best in May and October, corresponding to times of spring and fall migrations.

There are no indexes of abundance available for juvenile spotted seatrout. Juveniles were not abundant in any estuarine surveys, most of which used trawls and did not sample the shallow-water habitat preferred by spotted seatrout. Juveniles were most abundant in fall in Texas (Breuer 1973), in summer in Louisiana (Adkins et al. 1979; Juneau 1975), Mississippi (Waller and Sutter 1982), Florida (Jannke 1971), Georgia (Mahood 1975), and North Carolina (Spitsbergen and Wolff 1974; Purvis 1976; Wolff 1976).

Commercial landings data have been collected by the Federal government in each state since 1880. From 1880-1927 the survey was conducted on the average of once every five years, from 1927 to 1956 annual surveys were conducted, and since 1956 data has been collected on a monthly basis. It should be noted that commercial statistics, when biased at all, tend to be somewhat underestimated due to reporting failures inherent in their collection. Commercial landings may reflect true abundance trends, changes in effort, changes in gear restrictions, and/or closures of areas to commercial fishing.

Total commercial landings of spotted seatrout reached a peak at over 5,800 mt in 1945 (Table 12-9). From 1949-1964 landings fluctuated between 2,400 and 3,400 mt. Landings increased to over 4,000 mt in 1965, fluctuated between 3,200 and 4,400 mt until 1977, after which they declined to 1,605 mt in 1982.

Gulf of Mexico landings (west coast of Florida and Texas) have generally accounted for 60-80% of the total spotted seatrout commercial landings. Gulf landings exceeded 2,000 mt in most years from 1902 to 1976 (Table

12-9) (Figure 12-3), but have steadily declined since 1973. Highest landings in this region were reported from the west coast of Florida, followed by Texas and Louisiana. No landings were reported from Texas in 1982 since the Texas Legislature passed a bill prohibiting the commercial sale of Texas-caught spotted seatrout.

On the Atlantic coast commercial landings of spotted seatrout have been reported from Maryland to Florida (Table 12-9) (Figure 12-3). Total Atlantic coast landings were highest in 1945 (2,495 mt) and have since declined, fluctuating between 200 and 600 mt over the last two decades. Maryland landings of spotted seatrout never exceeded 12 metric tons and none have been reported since 1956. Virginia landings declined drastically from a high of 345 mt in 1944, and fluctuated from 1-89 mt from 1947-1982. North Carolina landings were highest in the 1930s (315-860 mt), declined to a low in 1968 (44 mt), increased in the early 1970s (147-304 mt), and have since declined to a low of 38 mt in 1982. Spotted seatrout landings in South Carolina peaked at 67 mt in 1945, ranged from 9 to 39 mt from 1950-1966, and fluctuated between 1 and 4 mt from 1976 to 1982. Georgia's landings peaked at 52 mt in 1936 and have not exceeded 7 mt over the past two decades except from 1972-1976 (12-14 mt). Highest commercial landings of spotted seatrout on the east coast were reported for Florida, peaking at 1,985 mt in 1945 and declining to 200-400 mt from 1957 to the present. Merriner (1980) noted that variability in annual reported catch is typical for spotted seatrout and seems to parallel the climatic conditions of the preceding spring and winter. For example, the cold winters of 1976 through 1978, during which inshore waters were less than 4.4°C for several weeks, were followed by declines in spotted seatrout landings, especially north of Florida (Table 12-9).

Recreational fishery statistics have not been routinely collected. Salt-water angling surveys were conducted at 5 year intervals from 1960 to 1970 (Clark 1962; Deuel and Clark 1968; Deuel 1973) and regional surveys were conducted in 1974 and 1975. The 1960-1970 surveys required fishermen to recall and report for a one-year period the number and average weight of each species caught. The 1974-1975 surveys reduced the recall period to two months, but still required fishermen to report the number and average weight of each species caught. The results of these surveys probably include some misidentified species and overestimates of the catch, although the magnitude of the overestimation is not known. Annual surveys were begun in 1979 which include a combination household survey and intercept survey (creel census). Although the results of the 1979 survey were published (Anonymous 1980), they are presently being corrected to correspond with 1980 census figures. The results of the 1980 survey (Anonymous 1984) are presented, but are not directly comparable with the 1965 and 1970 surveys due to the different methodologies (Table 12-10). All species of seatrouts were combined in the 1960 survey.

The recreational surveys indicate that the sport fishery catch probably exceeds the commercial harvest of spotted seatrout. The total estimated recreational catches of spotted seatrout for 1965 (49,052 mt), and 1970 (48,304 mt) were 17 times greater than the reported commercial landings for those years (2,924 and 2,778 mt, respectively). The estimated 1980

Table 12-9. Commercial landings (metric tons) of spotted seatrout by state, 1887-1983.

Year	Maryland	Virginia	North Carolina	South Carolina	Georgia	East Coast Florida	Atlantic Coast Total	West Coast Florida	Alabama	Mississippi	Louisiana	Texas	Gulf Coast Total	Total
1887	-	-	-	-	-	-	-	-	-	117	238	427	782	782
1888	-	-	-	-	-	-	-	232	103	127	237	395	1,094	1,094
1889	-	-	-	-	-	-	-	234	93	168	281	488	1,264	1,264
1890	-	-	-	-	-	-	-	250	95	169	297	508	1,319	1,319
1897	-	-	-	-	-	-	-	376	134	205	257	459	1,431	1,431
1902	-	-	-	-	-	-	-	843	117	215	489	507	2,171	2,171
1908	-	-	-	-	-	-	-	547	94	234	500	478	1,853	1,853
1918	-	-	-	-	-	-	-	768	63	161	540	431	2,263	2,263
1923	-	-	414	-	-	-	414	721	22	186	355	691	1,975	2,389
1927	-	-	-	-	-	-	-	1,171	54	274	373	771	2,643	2,643
1928	-	-	-	-	-	-	-	1,216	57	221	401	526	2,421	2,421
1929	-	-	-	-	-	-	-	1,334	58	174	233	534	2,333	2,333
1920	2	90	315	5	22	36	470	1,234	51	105	322	473	2,185	2,655
1931	*	27	503	7	5	264	806	1,095	49	98	348	492	2,082	2,888
1932	2	38	850	6	21	239	1,166	975	49	103	287	443	1,857	3,023
1933	4	74	-	-	-	-	78	-	-	-	-	-	-	78
1934	4	22	839	6	25	345	1,241	1,060	66	136	688	1,117	3,067	4,308
1935	2	51	-	-	-	-	53	-	-	-	-	-	-	53
1936	2	51	635	44	52	357	1,141	1,580	54	133	470	833	3,070	4,211
1937	3	64	409	37	29	299	841	1,254	76	169	448	956	2,903	3,744
1938	4	179	385	36	28	250	882	1,323	56	129	244	945	2,697	3,570
1939	3	74	605	18	31	270	1,001	1,359	58	113	325	673	2,528	3,529
1940	*	42	525	19	14	320	920	1,481	76	59	119	352	2,077	2,997
1941	12	42	-	-	-	-	54	-	-	-	-	-	-	54
1942	2	27	-	-	-	-	29	-	-	-	-	-	-	29
1944	2	345	-	-	-	-	347	-	-	-	-	-	-	347
1945	4	203	227	67	9	1,985	2,495	1,750	263	137	416	780	3,346	5,841
1946	1	195	-	-	-	-	196	-	-	-	-	-	-	196
1947	1	49	-	-	-	-	50	-	-	-	-	-	-	50
1948	-	40	-	-	-	-	40	*	180	72	228	269	749	789
1949	-	49	-	-	-	-	49	2,067	99	89	702	286	2,943	2,992
1950	-	42	219	12	15	661	949	1,458	39	52	400	265	2,214	3,163
1951	-	29	121	21	14	568	753	1,511	47	83	273	197	2,111	2,864
1952	-	33	292	39	11	675	1,053	1,575	66	240	263	217	2,371	3,421

Table 12-9. (continued)

Year	Maryland	Virginia	North Carolina	South Carolina	Georgia	East Coast Florida	Atlantic Coast Total	West Coast Florida	Alabama	Mississippi	Louisiana	Texas	Gulf Coast Total	Total
1953	-	39	287	23	8	252	609	1,185	60	408	243	265	2,161	2,770
1954	-	47	365	12	9	556	989	1,042	48	666	198	304	2,258	3,247
1955	-	47	201	63	20	413	744	928	71	917	231	382	2,529	3,273
1956	*	89	177	101	5	485	857	934	51	653	278	379	2,295	3,152
1957	-	54	262	25	2	402	745	1,162	33	124	291	408	2,018	2,763
1958	-	27	80	9	1	346	463	1,375	32	152	297	525	2,371	2,834
1959	-	63	176	17	*	348	504	1,281	51	146	313	503	2,294	2,898
1960	-	25	78	13	*	404	520	1,289	28	75	212	583	2,176	2,696
1961	-	33	95	25	1	340	494	1,194	52	109	281	507	2,143	2,637
1962	-	13	93	12	*	343	461	1,216	45	80	192	449	1,982	2,443
1963	-	12	105	22	2	363	504	1,197	60	67	209	540	2,073	2,577
1964	-	11	93	27	1	347	479	1,289	59	79	161	444	2,032	2,511
1965	-	18	79	16	3	309	426	1,605	73	80	208	533	2,498	2,924
1966	-	5	53	11	1	329	398	1,439	21	66	293	684	2,504	2,902
1967	-	2	56	1	3	282	334	1,196	41	78	282	690	2,286	2,620
1968	-	3	44	5	1	289	342	1,390	46	122	281	848	2,687	3,029
1969	-	8	86	4	1	308	307	1,097	44	100	327	532	2,100	2,407
1970	-	30	183	4	5	323	545	1,199	38	116	356	525	2,233	2,778
1971	-	20	153	11	7	222	413	889	62	178	509	674	2,313	2,726
1972	-	6	228	8	12	288	542	970	100	116	771	680	2,637	3,179
1973	-	4	277	3	12	302	598	1,009	159	177	1,146	893	3,374	3,972
1974	-	12	304	4	7	299	626	1,025	165	134	964	905	3,193	3,819
1975	-	33	287	8	14	243	585	984	47	119	860	823	2,833	3,418
1976	-	18	289	3	14	241	565	1,035	20	80	731	802	2,667	3,232
1977	-	2	147	*	7	224	380	726	10	67	818	611	1,905	2,285
1978	-	3	44	*	1	183	231	914	15	48	308	527	1,812	2,043
1979	-	2	48	1	2	216	269	946	34	50	362	467	1,859	2,128
1980	-	1	78	4	2	253	338	887	12	12	274	443	1,628	1,966
1981	-	2	51	*	*	334	387	895	12	18	266	293	1,484	1,871
1982	-	2	38	1	2	330	373	844	28	31	329	-	1,232	1,605
1983	-	2	75	2	3	206	288	816	27	24	514	-	1,381	1,587

- None reported

* Less than 1 mt




Figure 12-3. U.S. commercial landings of spotted seatrout by geographic region, 1950-1982.

Table 12-10. Spotted seatrout recreational catch and effort statistics from national saltwater angling surveys, 1965, 1970, and 1980.

Survey year	Catch		Number of successful anglers	Average weight		Catch per angler	
	Number	Weight Tbs. kg		Tbs.	kg	Number	Weight Tbs. kg
-----THOUSANDS-----							
1965 ¹							
South Atl.	12,559	18,209 8,267	365	1.45	0.66	34.41	49.89 22.65
Gulf	55,108	89,837 40,786	986	1.63	0.74	55.89	91.11 41.37
TOTAL	67,667	108,046 49,052	1,351	1.60	0.72	50.09	79.97 36.31
1970 ²							
South Atl.	13,992	25,040 11,368	432	1.79	0.81	32.39	57.96 26.31
Gulf	52,779	81,356 36,936	1,012	1.54	0.70	52.15	80.39 36.50
TOTAL	66,771	106,396 48,304	1,444	1.59	0.72	46,24	73,68 33.45
1980 ³							
South Atl.	1,978	2,180 989		1.10	.50		
Gulf	16,917	18,649 8,459		1.10	.50		
TOTAL	18,895	29,729 9,448		1.10	.50		

¹Deuel and Clark 1968.

²Deuel 1973.

³Anonymous in press.

recreational landings (9,448 mt) were 5 times greater than reported commercial landings (1,966 mt) for the same year. The recreational harvest of spotted seatrout in 1975 and 1976 in Texas represented about two thirds of the total catch of that species in Texas waters (Weaver 1977). Davis (1980) reported that 55% of the spotted seatrout landed in Everglades National Park, Florida from 1972 to 1977 were caught by recreational fishermen. Contrasting these data raises a question of data-set reliability (Merriner 1980). The Gulf of Mexico subregion accounted for 70% or more of the total recreational landings of spotted seatrout during all three survey years.

Although no commercial landings of spotted seatrout have been reported for Maryland in recent years, substantial quantities are caught by recreational fishermen. The 1979 and 1980 Maryland saltwater sport fishing surveys indicated that the catch of spotted seatrout in Maryland waters increased from 62 mt in 1979 to 241 mt in 1980. However, the 1980 results are probably inflated due to an error in coding some weakfish as spotted seatrout (Williams et al. 1982; Williams et al. 1983).

Age of recruitment to the recreational fishery is determined by minimum size limits in some states. In Alabama tournament rules prohibit entering fish <279 mm (11 in) which eliminates all but the faster growing age I+ fish (Tatum 1980). Age II+ fish make up about 18% of the total spotted seatrout catch and age III+ fish are the first age class fully vulnerable to the fishing tournament. Catch curve analyses from the Everglades National Park recreational fishery indicated that age at full recruitment may have changed from age III in 1959 (Stewart 1961) to age IV in 1979 (Rutherford 1982). This apparent shift in age at recruitment may be due to sample bias since Stewart (1961) made a special effort to collect small fish in the 1959 study and the minimum size limit in Florida is 30.5 cm TL (12 in).

Mortality rates were estimated for populations of spotted seatrout in Florida (Iversen and Moffett 1962; Rutherford et al. 1982) and Alabama (Tatum 1980) (Table 12-11). In Alabama mean annual mortality for fish older than age III, the first class fully vulnerable to the fishery, was estimated to be 49.8% and ranged from 36.2% in 1968 to 58.1% in 1975 (Tatum 1980). There was an inverse relationship between mean size and mean number of fish caught per tournament. Iversen and Moffett (1962), in a 4-month tagging study near Pine Island, Florida, estimated that natural mortality (M) was about four times larger than fishing mortality (F). Their estimates of F and M are probably too high, especially that of M (Perret et al. 1980). Males had higher rates of total annual mortality ($A=.82$) than females ($A=.77$) in Everglades National Park, Florida (Rutherford 1982). Total annual mortality rates of fully recruited spotted seatrout (combined sexes) in Everglades National Park increased slightly from 1959 to 1979, probably because of an increase in fishing mortality. Total annual mortality of females also increased slightly from 1959 to 1979 while it decreased slightly for males. Exploitation ratios were similar for all fish during both time periods and fishing mortality coefficients were higher for males than for females.

Winter cold shock of juveniles and adults has been cited as a primary factor in local and coastwide declines in spotted seatrout (Merriner 1980). Tabb (1966) noted that the spotted seatrout is very sensitive to changes in temperature. The death of large numbers of trout following severe cold spells was documented by Smith (1907), Hildebrand and Cable (1934), Storey and Gudger (1936), Gunter (1941), Gunter and Hildebrand (1951), Tabb (1958), Tabb and Manning (1961), and Moore (1976). There is usually only one kill per season in a particular area since once driven into deeper water the fish stay there for the remainder of the winter (Tabb 1958).

Catastrophic mortalities of spotted seatrout have also been attributed to hurricanes, excessive fresh water, red tide, and supersaturated dissolved oxygen conditions (Perret et al. 1980). Tabb and Manning (1961) reported a mortality following hurricane Donna (9 September 1960) which led to fish stranding and to turbulence which stirred the marl bottom of upper Florida Bay and packed fishes' gill chambers. Tabb (1966) suggested that lower salinities caused by run-off from tropical storms may cause mortality of young fish; however, he did not find dead fish to support his hypothesis. Springer and Woodburn (1960) listed spotted seatrout as one of the species killed by a red tide (*Gymnodinium breve*) in the Tampa Bay area in fall 1957. A phytoplankton bloom in Galveston Bay created supersaturated dissolved oxygen conditions and resulted in the formation of gas bubbles within the bloodstream and other body areas of spotted seatrout (Renfro 1963).

Yield per recruit models were generated for the Everglades National Park recreational fishery (Rutherford 1982). Separate models were generated for males and females because of differences in growth. The current yield of 249 g calculated for females was lower than that calculated for males (265 g) and was obtained at a lower fishing mortality rate. Maximum yield could be reached at the current level of fishing mortality for each sex by increasing the minimum size limit to 340 mm (age 5, 15.5 in TL) for males and 398 mm (age 5.5, 18 in TL) for females. Calculated yield per recruit was very similar for the recreational fisheries in 1959 and 1979. In both years the recreational fishery harvested females at slightly less than the maximum yield per recruit, while calculated yield of males was near maximum. Given the 300 mm (12 in) size limit, yield per recruit could have been maximized by slightly increasing the fishing mortality in both years.

12.4.3 Community Ecology

The spotted seatrout is essentially a non-migratory, euryhaline, estuarine species. Its entire life history is spent in the estuarine habitat, particularly the nontidal portions with extensive submerged vegetation where seasonal fluctuations in temperature and salinity rather than daily fluctuations are the controlling factors. Wide tolerance to changes in estuarine conditions has allowed spotted seatrout to occupy a niche that is intolerable to most marine predators and competitors (Tabb 1966). Although spotted seatrout are estuarine-dependent, they do move seaward through tidal inlets in response to environmental extremes (Section 12.3.5).

Table 12-11. Annual (A) and instantaneous (F,M,Z) mortality estimates, exploitation ratios ($E=F/Z$) and ratios of conditional fishing ($m=1-e^{-F}$) to conditional natural mortality ($n=1-e^{-M}$) for spotted seatrout.

Location	Sex	A	F	M	Z	E	m/n
Pine Island, Florida ¹	Combined	0.99	1.44	5.80	7.24		
Alabama ²	Combined	0.50*					
Everglades National Park, Florida ³	Combined						
	1959	0.72	0.83	0.43	1.26	0.66	1.61
	1979	0.77	1.03	0.45	1.48	0.69	1.77
	<u>Males</u>						
	1959	0.85	1.36	0.54	1.90	0.72	1.78
	1979	0.82	1.26	0.46	1.72	0.73	1.94
	<u>Females</u>						
	1959	0.66	0.68	0.39	1.07	0.63	1.40
	1979	0.75	0.88	0.50	1.38	0.64	1.50

¹Iversen and Moffett 1962

²Tatum 1980

³Rutherford 1982

*Mean mortality rate 1968-1975 (range: 0.36-0.58).

Klima and Tabb (1959) noted that spotted seatrout on the east coast of Florida attained a larger mean length at each age compared with those on the northwest coast and suggested that this might be due to environmental differences between the areas. In Apalachee Bay (northwest Florida) there is a scarcity of protected calm water areas and the grass flats are composed of essentially marine species (turtle grass and manatee grass). The Indian River area (east-central Florida) consists of large areas of shallow, quiet, brackish waters supporting dense stands of shoal grass and a distinctive brackish water fauna. Nine species of spotted seatrout predators and numerous competitors were listed for Apalachee Bay (Table 12-4) compared with only five species of predators for the east coast. Klima and Tabb (1959) suggested that the smaller number of predators in the Indian River environment is probably a result of the inability of many marine species to invade low salinity waters for any length of time. Spotted seatrout have successfully invaded the rich feeding grounds of the euryhaline herbivores and under ideal conditions may be the top carnivore (Tabb 1958).

12.5 Exploitation

12.5.1 Commercial Exploitation

The commercial fisheries for spotted seatrout were reviewed by Goode (1887), Brice et al. (1898), Tabb (1958), Klima and Tabb (1959), Moffett (1961), Anderson and Gehringer (1965), Merriner (1980), and Perret et al. (1980).

12.5.1.1 Fishing Equipment

The principal commercial methods used to harvest spotted seatrout include various types of gill nets (runaround, stake, anchor, set, and drift), haul seines, pound nets, hand lines, troll and trot lines, trammel nets, and otter trawls (fish and shrimp). Runaround gill nets, haul seines, and anchor, set or stake gill nets accounted for 86% of the Atlantic coast catch, whereas over 84% of the landings in the Gulf of Mexico came from runaround gill nets, trammel nets, and trot lines (Tables 12-12 and 12-13).

Gear use varies among states and is partly a function of gear efficiency in different areas, but probably is more a function of state or local laws (Perret et al. 1980). For example, no gill netting is permitted in Georgia. The commercial landings data indicate that the majority of spotted seatrout landed in Georgia are caught by hook and line. In South Carolina they are caught in drift gill nets, hand lines, and incidentally in shrimp trawls. Prior to 1971, however, most were caught in haul seines. Highest landings in Virginia and North Carolina are from haul seines followed by drift gill nets in Virginia and anchor gill nets in North Carolina. In Florida, spotted seatrout are mainly caught by runaround gill nets.

Seasonal changes in use of commercial gear types occur in Florida and North Carolina. Trammel nets and haul seines are used primarily during the winter months in the rivers of northwest Florida for mullet, spotted seatrout and red drum (Klima and Tabb 1959). Hook and line fishing is

Table 12-12. Atlantic coast catch of spotted seatrout by gear type, 1976 (metric tons).

Gear Type	Virginia	North Carolina	South Carolina	Georgia	Florida	Total	%
Runaround gill net	-	22.5	-	-	206.7	229.2	40.6
Haul seine	8.6	165.3	-	-	6.0	179.9	31.9
Anchor, set or stake gill net	1.7	73.3	-	0.9	0.3	76.2	13.5
Pound net	1.4	25.7	-	-	-	27.1	4.8
Hand line	-	-	0.3	2.2	19.6	22.2	3.9
Drift gill net	5.9	2.2	2.1	1.2	-	11.4	2.0
Shrimp otter trawl	-	0.3	0.2	9.3	1.3	11.0	1.9
Trammel net	-	-	-	-	4.9	4.9	0.9
Troll line	-	-	-	-	2.5	2.5	0.4
Fish otter trawl	0.2	-	-	-	-	0.2	0.0
						564.6	99.9

Source: Fishery Statistics of the United States, 1976, National Marine Fisheries Service

- None reported

Table 12-13. Gulf of Mexico catch of spotted seatrout by gear type, 1976 (metric tons).

Gear type	Florida	Alabama	Mississippi	Louisiana	Texas	Total	%
Runaround gill net		654.1	-	69.4	549.3	-	1,272.8 47.6
Trammel net		136.2	17.5	.1	150.6	290.6	595.0 22.3
Long (trot) line		-	-	-	0.2	387.3	387.5 14.5
Haul seine		103.2	-	-	2.1	92.4	197.7 7.4
Hand line		104.4	1.9	2.0	6.2	6.8	121.3 4.5
Anchor, set or stake gill net		-	-	-	22.8	25.6	48.4 1.8
Troll line		26.7	-	-	-	-	26.7 1.0
Drift gill net		-	-	9.1	-	-	9.1 0.3
Shrimp otter trawl		6.6	0.5	-	0.2	0.3	7.6 0.3
Fish otter trawl		4.8	-	-	-	-	4.8 0.2
Purse seine		-	-	0.9	-	-	0.9 0.0
						2,671.8	99.9

Source: Fishery Statistics of the United States, 1976, National Marine Fisheries Service.

- None reported

productive throughout most of the year in west Florida, whereas trolling is usually best in the fall. According to local fishermen, the best gill and trammel net fishing is from mid-November to mid-February when fish congregate in deep holes, and also in the spring (Moffett 1961). In North Carolina, spotted seatrout are caught mainly by long haul seines and pound nets in spring and summer, long haul seines and gill nets in fall, and trawls and gill nets in winter.

12.5.1.2 Areas Fished

The best fishing areas for spotted seatrout in Florida and throughout its range are shallow brackish bays, lagoons, mangrove-bordered estuaries with abundant submerged aquatic vegetation and adjacent deep holes or channels (Tabb 1958, 1960, 1966). In Chesapeake Bay largest catches of spotted seatrout were made in the lower bay, followed by the Rappahannock and York Rivers (Hildebrand and Schroeder 1928). Largest catches of spotted seatrout in North Carolina are made in Pamlico Sound. Commercial landings statistics indicate that the majority of spotted seatrout catches are made in the estuaries along the Atlantic Coast (82-99% in 1982) (Table 12-14).

12.5.1.3 Fishing Seasons

In Chesapeake Bay spotted seatrout are caught from March until December with periods of peak abundance from March to May, and September to November. Spotted seatrout are caught year-round within estuaries from North Carolina southward and offshore of North Carolina in winter. Largest catches are made in the fall (October-December) in North Carolina. On the east coast of Florida largest catches are made during winter when spotted seatrout are concentrated in channels and deep holes in estuaries and also in spring at spawning time (Tabb 1960; Anderson and Gehringer 1965).

12.5.1.4 Fishing Operations and Results

Limited catch per unit of effort data are available for spotted seatrout commercial fisheries in the Gulf of Mexico. Davis (1980) presented commercial and recreational effort data for the Everglades National Park fisheries, 1972-77 and compared mean weights and harvests. The number of gill net sets fluctuated over the 6-year period and varied between areas within the Park. Man-hours of commercial line fishing declined in all areas. The commercial catch accounted for 45% of the total Park spotted seatrout harvest.

Matlock et al. (1979) compared catch rates of spotted seatrout in areas open and closed to commercial netting. The overall mean catch rate from areas closed to commercial netting was approximately twice as high as that from open areas. There was no difference in mean size of fish between the areas.

Klima and Tabb (1959) reported on gear selectivity in the spotted seatrout fishery in northwest Florida. The sizes of fish caught by commercial anglers were influenced by the size of the hook and size of

Table 12-14. Commercial catch of spotted seatrout (kg) by water area for the Atlantic coast (excluding Florida) and percent (%) caught within estuaries.

Year	Virginia		%	North Carolina		%	South Carolina		%	Georgia		Total %	
	Ocean	Estuary		Ocean	Estuary		Ocean	Estuary					
1972	544	52,617	99	36,015	194,315	84	2,592	5,544	68	5,734	6,010	51	85
1973	544	3,765	87	84,368	192,779	70	1,894	729	28	6,672	5,476	45	68
1974	2,268	9,616	81	38,617	265,112	87	577	3,444	86	2,889	4,429	61	86
1975	5,216	27,669	84	37,042	249,902	87	1,846	5,852	76	6,571	7,388	53	85
1976	1,315	16,375	93	11,835	277,251	96	188	2,442	93	6,715	6,858	51	94
1977	318	1,406	82	6,297	140,399	96	61	176	74	4,903	1,903	28	93
1978	454	2,313	82	8,326	35,310	81	26	28	52	292	828	74	81
1979	318	1,270	80	9,373	38,270	80	0	2,092	100	79	2,183	97	82
1980	91	363	80	17,214	60,502	78	2,638	1,053	28	0	1,928	100	76
1981	45	1,769	97	3,502	47,892	93	38	84	69	5	281	98	93
1982	45	1,497	97	5,249	32,784	86	162	720	82	1	2,264	99	87

bait. Spotted seatrout caught by hook and line had a smaller average size (28.5 cm SL) than those caught in gill nets (30.5 cm SL) or seines (34.5 cm SL).

An evaluation of monofilament and multifilament gill nets with 1-5/8", 1-7/8", and 2" bar mesh in Louisiana indicated that monofilament nets with 1-5/8" bar mesh were most efficient in capturing spotted seatrout (Adkins and Bourgeois 1982). The size range of 482 fish was 345-451 mm TL and the average size was 410 mm TL. Hein and Shepard (1980) reported that the mean length of all fish captured in 3 to 3-1/4" stretch monofilament gill nets in Louisiana was 377 mm TL, with a mean weight of 544 g. The mean size for females was 391 mm TL (611 g) and for males was 354 mm TL (440 g). In Georgia monofilament gill nets of 2-7/8" stretched mesh and caught spotted seatrout 238-633 mm TL, with 92% ranging from 303-428 mm TL (Mahood 1975). Tabb (1960) reported an average size of 335 mm SL and 0.6 kg for spotted seatrout taken in 3-1/8" stretched mesh gill nets in Florida.

12.5.1.5 Incidental Catches

Spotted seatrout apparently do not contribute significantly to the incidental finfish catches in the South Atlantic or Gulf coast shrimp fisheries. Spotted seatrout were not reported in a study of the scrap fishery of North Carolina (Wolff 1972), in a shrimp trawling investigation along the coast of South Carolina or Florida (Anderson 1968) or in a study of shrimp fishing in Georgia's close inshore waters (Knowlton 1972). Anderson (1968) reported total catches of 54 fish from Georgia outside waters (coast to 11.1 km offshore) and 224 fish from Georgia inside waters (rivers, creeks, and sounds), both less than 0.05% of the total number of finfish caught during shrimp trawling, 1931-1935. Bearden (1969) noted that commercial shrimp trawling efforts had little effect on spotted seatrout populations in South Carolina because juveniles are found in the inshore, estuarine areas (off limits to shrimp trawling) and adults have the mobility to evade trawl gear. Spotted seatrout comprised only 0.02% by number (45,000/yr) and 0.08% by weight (5,800 kg/yr) in the South Carolina shrimp fishery, 1974-75 (Keiser 1976). Landings statistics from the National Marine Fisheries Service, however, indicated that sizeable quantities of spotted seatrout were harvested in shrimp trawls during the mid-1960s and early 1970s. Higgins and Pearson (1928) reported that only a small percentage by number (1.8-14.0%) of spotted seatrout in North Carolina long haul seine catches were unmarketable.

12.5.2 Recreational Exploitation

Aspects of the recreational fisheries for spotted seatrout were reviewed by Tabb (1960), Anderson and Gehringer (1965), Higman (1967), Freeman and Walford (1974, 1976a,b,c,d), Davis (1980), Merriner (1980), Perret et al. (1980) and Brown (1981).

12.5.2.1 Fishing Equipment

Spotted seatrout are caught by anglers while bottom fishing, chumming, live lining, jigging and casting from shore, and trolling from boats

(Freeman and Walford 1974, 1976a,b,c,d). The salt-water angling surveys indicate that the principal mode of fishing for spotted seatrout is from private or rented boats (Deuel and Clark 1968; Deuel 1973; Anonymous 1984) (Table 12-15). Principal baits include shrimp, mullet, soft or shedder crabs, silversides and killifish. Lures such as plugs, weighted bucktails, jigs, spoons, spinners and streamer flies are also used (Freeman and Walford 1974, 1976a,b,c,d).

In Chesapeake Bay bait fishing, casting, and trolling are the most often employed fishing techniques (Brown 1981). The most effective bait is peeler crabs and the most successful lures are stingray grubs, bucktails, and mirrolures. Bait is used mainly in spring and summer, and lures in fall. The best all-round outfit for spotted seatrout is a light spinning outfit with six-foot rod calibrated for 1/4 to 1/2-oz lures with a small reel with 8 or 10-lb mono (Osborne 1981). Lures are apparently more popular in North Carolina waters and more successful in catching citation size fish (Brown 1981).

In South Carolina anglers drift or anchor over deep holes, cast along shell banks or near pilings, troll, or surf and pier fish for spotted seatrout. Live shrimp is the most popular bait, but dead shrimp or mud minnows are also used. Many fishermen prefer to use lures such as the bucktail and stingray grub (Cupka 1972).

Spotted seatrout is the most popular sportfish in coastal Georgia and most are caught on live shrimp (Anonymous 1983b). A recent Georgia tagging study indicated that 64% of all recaptured spotted seatrout were taken using live shrimp, 27% using artificial lures, and the remainder using dead shrimp, cut bait, minnows, and fiddler crabs. During cooler weather (mid-November through March) artificial lures work just as well and often better in upper rivers and creeks. When using artificial lures, fishing tackle usually consists of light spinning or spin cast reels with rods 6 feet long or longer (12 lb line or lighter). Spotted seatrout fishing is generally less productive for two days before and after the new and full moon phases because high tidal ranges cause muddy waters (Anonymous 1983b).

Spotted seatrout fishing in Florida includes bridge, skiff, and shoreline fishing. Live bait, including shrimp, sailors choice, pinfish, mullet, and needlefish, is generally used with greater success than lures, although experienced anglers are successful using the latter (Tabb 1960).

12.5.2.2 Areas Fished

The 1965, 1970, and 1980 saltwater angling surveys indicated that the majority of spotted seatrout were caught in sounds, rivers, and bays (Deuel and Clark 1968; Deuel 1973; Anonymous 1984) (Table 12-15). The best spotted seatrout fishing in Chesapeake Bay in summer occurs in areas with abundant grass beds, particularly on the bayside of the Eastern Shore (Brown 1981). In the fall spotted seatrout catches are best in areas with adjacent deep water such as Smith Island, Magothy Bay, Lynnhaven River, Rudee Inlet, and the Chesapeake Bay Bridge Tunnel.

Table 12-15. Number of spotted seatrout caught by U.S. anglers in each region in 1965, 1970, and 1980 by principal area and method of fishing.

Survey year	Region	Principal area of fishing		Principal method of fishing					Beach or bank
		Ocean	Sounds rivers and bays	Private or rented boat	Party or charter boat	Bridge pier or jetty			
----- THOUSANDS -----									
1965	South Atlantic	2,343	10,216	10,173	180	1,412		794	
	Gulf of Mexico	8,597	46,511	41,245	2,448	8,325		3,090	
1970	South Atlantic	4,374	9,618	7,686	2,187	3,433		686	
	Gulf of Mexico	16,322	36,457	38,499	2,818	5,694		5,768	
1980	South Atlantic	72	1,675	1,682	12	193		91	
	Gulf of Mexico	4,169	8,802	13,990	986	1,029		912	

¹Deuel and Clark 1968

²Deuel 1973

³Anonymous 1984

In North Carolina spotted seatrout are most frequently caught around islands and points, over grass flats and oyster bars, at creek mouths, at jogs and bends in channels, near jetties and pilings, in marshes crisscrossed by creeks and guts, and along steeply cut banks (Osborne 1981). In late October and November spotted seatrout can be found in the surf of the Outer Banks from Corolla to Portsmouth Island (Randolph 1983). Most North Carolina citation catches have come from Pamlico Sound and Oregon Inlet (Brown 1981). The most popular areas for spotted seatrout fishing in South Carolina include Murrells Inlet, North Edisto River, Wando River and the numerous estuarine areas and tidal creeks in the southern part of the state (Cupka 1972). In Georgia they are found concentrated in areas with large quantities of dead shell, or adjacent to live oyster beds. Schools may be found in the surf zone along the beach during the warmer months. Spotted seatrout are also caught at night from lighted piers and docks. (Anonymous 1983b). On the east coast of Florida the greatest numbers of spotted seatrout are landed in the Indian River lagoon system (Tabb 1960).

12.5.2.3 Fishing Seasons

The catch per unit effort for spotted seatrout in Maryland was highest in September-October (Williams et al. 1982; Williams et al. 1983). In Chesapeake Bay spotted seatrout are caught by anglers from May-October; the best month is October, followed by May (Brown 1981). In North Carolina best fishing begins in July, peaks in October, and continues to December or well into January in a mild year. The prime fishing months for spotted seatrout in South Carolina are September-December and May-June (Cupka 1972). Spotted seatrout fishing takes place year-round in Florida waters (Tabb 1960). Anderson and Gehringer (1965) reported highest recreational catches in spring in the Cape Canaveral area but did not sample during winter.

12.5.2.4 Fishing operations and results

Recreational catch per unit effort data for spotted seatrout are not available for the Atlantic coast. Catch rates for Everglades National Park, Florida, from 1958 through 1978 were presented by Higman (1967) and Davis (1980). Catch per unit effort data for the Gulf states was summarized by Perret et al. (1980).

12.6 Social and Economic Implications

12.6.1 Values

Spotted seatrout contributed more to the total value of U.S. sciaenid landings between 1960 and 1974 than any other species (Cato 1981). Total value of spotted seatrout landings have ranked third behind croaker and weakfish since 1977 (\$3.0 million in 1982).

The total value of Gulf of Mexico landings generally increased from 1950 to 1982 (Figure 12-4). Atlantic coast values of spotted seatrout landings have fluctuated, but increased from 1979 to 1982. Overall price movements have been fairly consistent in both regions with Gulf of Mexico prices usually slightly below prices in the South Atlantic prior




Figure 12-4. Dockside value of U.S. commercial landings for spotted seatrout, 1950-1982.

to 1975 (Table 12-16). Adjusting prices for inflation indicates that the real (deflated) price of spotted seatrout declined from 1967 to 1977 along the Atlantic coast. Gulf of Mexico prices have increased since 1974 (Cato 1981).

Cato (1981) analyzed spotted seatrout monthly price movements for Florida. Monthly prices were lowest in the winter months when landings were heaviest. The effect on price of a one-million-lb (454 mt) increase in monthly Florida landings was a \$.086 decrease, or slightly less than a 1-cent decrease for each increase of one hundred thousand pounds (45.4 mt). A 1% increase in landings explains a .04% decrease in price. Other factors important in explaining spotted seatrout price variation were previous months' landings, total personal income, and quantities of other species landed thought to substitute in the market for spotted seatrout. A one-million-lb (454 mt) increase in the landings of mullet, sheepshead, flounder, croaker, and red drum in Florida was seen to explain almost one-half of a 1-cent decrease in spotted seatrout prices. This indicates that these fish may be good substitutes for each other in the marketplace.

12.6.2 Employment

There are no data available on employment in the various spotted seatrout fisheries. Tabb (1960) stated that commercial fishing effort was declining along Florida's east coast because of closed commercial netting in some inshore waters and due to rapid urbanization and industrialization of this area.

12.6.3 Participation

The spotted seatrout is one of the most sought after and most often caught species of sportfish in its range (Tabb 1960; Cupka 1972). Its wide geographic range, desirable food value, and angling qualities contribute to this popularity. Participants in the spotted seatrout fishery include commercial fishermen, processors and dealers, food consumers, and recreational fishermen.

Few data are available on commercial fishing investment, total effort, efficiency, productivity, and costs for the spotted seatrout fishery, which is a mixed species fishery. Anderson and McNutt (1973) reported that spotted seatrout and red drum represented 8% of \$20,000 in total returns for a small boat gill net fisherman on Florida's west coast.

Sport fishermen in the Indian River area of Florida were divided into three categories by Tabb (1960) based on disposition of the catch: fishermen who fish for recreation and home consumption only; those who consider themselves sportsmen, but who market some fish to defray trip expenses; and those who fish for sport, but who always market their catch. The South Carolina gill net fishery is primarily a noncommercial fishery; fishermen utilize small nets (<30 m), to supply fish for home consumption. Only 6% of the gill net fishermen who fished in 1978 sold a portion of their catch. Spotted seatrout comprised about 4% of the catch (7,500 kg) (Moore 1980). Hammond and Cupka (1977) made an

Table 12-16. Spotted seatrout ex-vessel prices, 1967-1977 (cents per pound).

Year	Atlantic Coast		Gulf of Mexico	
	Reported ¹	Adjusted ²	Reported ¹	Adjusted ²
1967	28.8	28.8	24.9	24.9
1968	30.1	29.4	26.0	25.4
1969	32.6	30.6	27.9	26.2
1970	30.2	27.4	27.9	25.3
1971	31.6	27.7	27.6	24.2
1972	34.6	29.1	29.6	24.9
1973	33.8	25.1	33.7	25.0
1974	35.3	22.0	33.7	21.0
1975	37.6	21.5	39.2	22.4
1976	39.6	21.6	45.2	24.7
1977	43.2	22.1	50.7	26.1

¹Value divided by landings

²Reported price adjusted by wholesale price index

economic evaluation of the South Carolina pier fishery and found that spotted seatrout was a relatively minor component in this fishery (<1% of total catch).

12.6.4 Processors and Product Forms

Spotted seatrout are marketed primarily along the coastal states of the Gulf of Mexico and in adjoining states (Cato 1981). In 1956 approximately 58% of all the spotted seatrout landed in the states of North Carolina, South Carolina, Georgia, Alabama, Mississippi, Texas, Louisiana, and Florida came from Florida waters (Rosen and Ellis 1958). A survey of 142 retail and wholesale markets in Georgia, Florida, and Alabama revealed that spotted seatrout sales were fairly consistent all year. Spotted seatrout were offered by 96% of the markets and average market sales per week ranged from 175 to 227 kg (384-501 lb). The predominant sales form was fresh and the average price per pound (per 0.45 kg) paid during abundant supply seasons was \$1.01 (Anonymous 1979). Most seafood retailers in South Carolina reported selling about 11 kg of spotted seatrout or less per year although a few reported as much as 1,360 to 1,800 kg per year. Hook and line and gill net fishermen are the primary source of supply (Smith and Moore 1979). Perret et al. (1980) reported that virtually all of the commercial landings in the Gulf are sold in local markets as fresh in the round or gutted. A small percentage is sold as frozen and gutted or as fresh or frozen fillets.

12.6.5 Import/Export

Perret et al. (1980) stated that imports of spotted seatrout from Mexico are substantial, and have occasionally exceeded 454 mt (Table 12-17). These imports have an impact in Texas, Louisiana, Oklahoma, and perhaps other markets supplied from Texas and Louisiana landings. The net impact of imports is not known.

12.6.6 Gear Conflicts

Gear conflicts may occur between the long haul seine fishermen and the pound net, crab and eel pot fishermen in North Carolina. Abandoned, broken-off pound net stakes and pound net stakes left in place from season to season exclude long haulers from large areas, especially in Core Sound. A very large increase in the number of crab and eel pot fishermen has resulted in ever increasing friction with haul seiners, who cannot haul in areas filled with pots. Potters are mainly interested in shoal waters, which long haulers need only to bunt or harden up their seine (DeVries 1981).

12.6.7 Commercial-Recreational Conflicts

In Florida there is some contention that bait shrimp trawlers and commercial netting (gill, trammel, and seines) negatively impact spotted seatrout fishing by killing vast quantities of juveniles in the estuaries as well as damaging seagrass beds (Futch 1970). There has been legislation introduced in Florida to ban gill netting for spotted seatrout; however, none of it has passed. Commercial-recreational

Table 12-17. Imports of spotted seatrout, 1950-1977. (from Perret et al. 1980)

Year	Metric tons
1950	562.7
1951	465.4
1952	553.2
1953	558.4
1954	593.1
1955	589.1
1956	642.8
1957	726.5
1958	714.1
1959	773.4
1960	679.0
1961	121.7
1962	146.5
1963	164.8
1964	153.8
1965	121.2
1966	103.9
1967	58.5
1968	62.4
1969	375.1
1970	589.1
1971	466.0
1972	317.6
1973	317.5
1974	429.3
1975	379.7
1976	365.7
1977	631.1

Source: U. S. Bureau of Customs Records Transcribed by National Marine Fisheries Personnel.

conflicts in Everglades National Park were discussed by Davis (1982). The National Park Service has imposed bag limits on recreational fishermen and proposed to eliminate commercial fishing in the Park by December 31, 1985. In North Carolina there is a growing conflict between recreational anglers and long haul fishermen (DeVries 1981). Conflicts and controversies in the Texas spotted seatrout fisheries were reviewed by Heffernan and Kemp (1982) and Matlock (1982). Regulations to close Texas bays to commercial fishing were adopted in the early 1900s and legislative action was taken from the 1930s to the 1970s to reduce commercial fishing pressure on the stocks, which included size limits, opened and closed waters, and gear restrictions. In 1981 legislation was passed which prohibited the sale of Texas-caught spotted seatrout. In January 1983 a task force of administrators and biologists from the Louisiana Department of Wildlife and Fisheries issued a report recommending creation of a Finfish Research/Management Section. The task force was created in late 1982 as the direct result of a controversy between commercial and recreational fishermen over laws governing spotted seatrout and red drum.

12.7 Management and Protection

12.7.1 Regulatory Measures

The fisheries for spotted seatrout have been conducted almost entirely within the internal waters of the states and in the territorial sea which extends 5.6 km (3 n mi) offshore on the Atlantic coast. Therefore, management has been by individual state regulation. Regulations and methods of promulgating them vary among states and are summarized in Table 12-18. The only regulations specifically dealing with spotted seatrout are minimum size limits of 23 cm (9 in) in Maryland and 30 cm (12 in) in Florida.

The Atlantic States Marine Fisheries Commission (ASMFC) administers a cooperative program with the National Marine Fisheries Service (NMFS) entitled the Interstate Fisheries Management Program (ISFMP). This program provides funding to the Atlantic coastal states to coordinate interjurisdictional fisheries management and develop fishery management plans (FMPs) for species occurring in the territorial sea. Plans for coastal migratory species such as Atlantic menhaden, summer flounder, and striped bass have been developed under the ASMFC program and several states have implemented regulations in compliance with these plans.

The Magnuson Fishery Conservation and Management Act (MFCMA) provides for the conservation and exclusive management by the Federal government of all fishery resources within the United States Fishery Conservation Zone (FCZ). The FCZ extends from the territorial sea to 370 km (200 n mi) from shore. Fishery management in the FCZ is based on fishery management plans developed by regional Fishery Management Councils (FMC). Spotted seatrout rarely occur in the FCZ, except off North Carolina in winter.

The National Park Service retains the authority to manage fish primarily through the establishment of coastal and nearshore national parks and

Table 12-18. Synoptic overview of present state management systems.

	Rhode Island	Connecticut
Administrative organization	Rhode Island Department of Environmental Management	Connecticut Department of Environmental Protection
Legislative organization	Rhode Island Marine Fisheries Council	Connecticut Commissioner Environmental Protection
Licenses	Commercial	Commercial
Size restrictions	None	None
Limits	None	None
Gear restrictions	None	None
Conservation regulations	None	None

Table 12-18. Continued

New York	
Administrative organization	New York State Department of Environmental Conservation
Legislative organization	New York Fish and Game Laws, Article 13 Marine and Coastal Resources
Licenses	Commercial non-resident beam and otter trawl
Size restrictions	None
Limits	None
Gear restrictions	Trawl prohibited from Great South Bay, Moriches Bay, Shinnecock Bay; seasonally in Peconic Bays. Gill nets restricted from Peconic Bays; haul seines limited in lengths in these same bays and cannot be fished from midnight Thursday to 6:00 p.m. Sunday. Nets and trawls may not be set in western Long Island Sound Apr. 1 - Nov. 1. Gill nets prohibited in central and western Long Island Sound.
Conservation regulations	None

Table 12-18. Continued

	New Jersey	Delaware
Administrative organization	New Jersey Department of Environmental Protection, Division of Fish, Game and Wildlife, Marine Fisheries Administration, Bureau of Marine Fisheries	Division of Fish and Wildlife Department of Natural Resources and Environmental Control
Legislative organization	New Jersey Statutes, Title 23, Chapter 28	Delaware State Legislature
Licenses	Fyke nets - \$1, \$4, \$30 Haul seines - \$25 Bait seines - \$3 (50' - 150') Gill nets - anchored - \$13 drift - \$20 run around - \$20 Pound nets - \$ 25 - \$ 50 - \$100 Otter trawl - \$100 Beam trawl - \$100 Purse seine - \$100	None
Size restrictions	None	None
Limits	None	None
Gear restrictions	Trawls and purse seines restricted from within 2 miles of coastline. Seasons for gill nets, fyke nets, haul seines.	Trawls prohibited in Delaware Bay. Gill nets, fyke nets and seines allowed. Purse seines prohibited within 3 miles of coast.
Conservation regulations	None	None

Table 12-18. Continued

	Maryland	Virginia
Administrative organization	Maryland Department of Natural Resources	Virginia Marine Resources Commission
Legislative organization	Natural Resources Article, Annotated Code of Maryland Title 4, Subtitle 1, Title 08, Subtitle 02, Chapter 05 Fish	Marine Resources of the Commonwealth Code of Virginia of 1950, Title 28.1
Licenses	Otter trawl - \$100 Beam trawl - \$100 Fyke or hoop nets - \$50 Gill nets- <200 yds \$100 >200 yds \$200 Chesapeake Bay sport fishing license \$5	Commercial
Size restrictions	9" Minimum	None
Limits	None	None
Gear restrictions	Trawling prohibited within 1 mile of Maryland shoreline in Atlantic Ocean. Numerous gear and area restrictions.	Trawling prohibited in Chesapeake Bay. Pound net mesh <2" (s.m.) and haul seine mesh >3" mesh (s.m.) prohibited. No trot or long lining along the ocean side of Eastern shore. Encircling gill nets prohibited. 4½" minimum net mesh in Virginia waters.
Conservation regulations	Secretary of Natural Resources has authority to adopt rules and regulations relating to taking, possession, transportation, exporting, processing, sale or shipment necessary to conservation.	None

Table 12-18. Continued

North Carolina	
Administrative organization	North Carolina Department of Natural Resources and Community Development, Division of Marine Fisheries
Legislative organization	North Carolina Administrative Code, Title 15, Chapter 3.
Licenses	<p>Vessels without motors, any length, when used with other licensed vessel - no license</p> <p>Vessels, not over 18' - \$1/foot</p> <p>Vessels, over 18' to 38' - \$1.50/foot</p> <p>Vessels, over 38' - \$3/foot</p> <p>Non-resident vessels - \$200 in addition to above fee requirement</p> <p>Finfish processor - \$100</p> <p>Unprocessed finfish dealer - \$50</p>
Size restrictions	None
Limits	None
Gear restrictions	Trawling for finfish prohibited in internal coastal waters. No purse seine for food fish. Many specific net regulations for areas and seasons.
Conservation regulations	Secretary, acting upon advise of Director of Marine Fisheries, may close area to trawling if in coastal fishing waters, samples become composed primarily of juvenile finfish of major economic important.

Table 12-18. Continued

	South Carolina	Georgia
Administrative organization	South Carolina Wildlife and Marine Resources	Georgia Department of Natural Resources
Legislative organization	Section 50-5-20	Georgia Code 27-4-110
Licenses	Land and sell \$25 Commercial boat license $\leq 18'$ - \$20 $> 18'$ - \$25 Gill nets haul seines - \$10/100 yds	Commercial fishing license (personal) - \$10.25 for any sales of catch Nontrawler license $\leq 18'$ - \$5 $> 18'$ - \$5 + \$.50/foot Trawler license - \$50 for 18' + \$3/additional foot No license for seines $< 100'$ unless catch is sold.
Size restriction	None	None
Limits	None	None
Gear restrictions	Seine mesh $< 2\frac{1}{2}"$ prohibited Purse seining for food fish permitted in ocean greater than 300 yds from beach.	Gill netting prohibited in Georgia waters. Seine mesh restrictions: minimum of $1\frac{1}{4}"$ for seines less than 100'; minimum mesh size of $2\frac{1}{2}"$ (stretched mesh) for 100' - 300' maximum length.
Conservation regulations	None	None

Table 12-18. Continued

Florida	
Administrative organization	Marine Fisheries Commission
Legislative organization	Chapter 370, Florida Statutes; additional 220 state laws that apply on a local level; all local laws will become Rules of the Marine Fisheries Commission by July 1, 1985.
Licenses	License to sell: Resident - \$25 annually Non-resident - \$100 annually Alien - \$150 annually Wholesale seafood dealer Resident - \$300 annually Non-resident - \$500 annually Alien - \$750 annually Retail seafood dealer Resident - \$25 annually Non-resident - \$200 annually Alien - \$250 annually
Size restrictions	12" FL minimum except in Franklin and Wakulla counties
Limits	None
Gear restrictions	Purse seining and stop netting prohibited. Numerous local gear and area restrictions.
Conservation regulations	None

national monuments such as Everglades National Park in Florida.

12.7.2 Habitat Protection

The spotted seatrout is essentially a nonmigratory estuarine species, except perhaps at the northern extreme of its range. Nearly the entire commercial and recreational catch of spotted seatrout comes from estuaries (Tables 12-14 and 12-15). The habitat value of saltmarshes, mangroves, and seagrasses for aquatic organisms, including spotted seatrout was discussed by Thayer et al. (1978). Man's activities in these areas may negatively affect the suitability of the habitat for spotted seatrout and thereby reduce the natural production of this species (Merriner 1980).

Estuarine habitats have deteriorated rapidly since approximately 1940, mostly as a result of industrial and human population growth. The National Estuary Study, completed in 1970, indicated that 73% of the Nation's estuaries had been moderately or severely degraded (Gusey 1978, 1981). Damage and/or destruction of estuaries has largely been by dredging and filling for waterfront property, dredging of navigation channels, construction of causeways and bridges, installation of ports and marinas, alteration of freshwater flow, and pollution. Unfortunately the effects of habitat alterations have rarely been quantified.

The association of juvenile and adult spotted seatrout with seagrass beds is well documented (Pearson 1929; Miles 1950; Moody 1950; Reid 1954; Tabb 1958). Seagrass beds along the coast of Mississippi were virtually destroyed during Hurricane Camille in August 1969 (Lorio and Perret 1980); however, it is not known what effects this had on spotted seatrout populations in Mississippi Sound. An unprecedented decline in submerged aquatic vegetation has occurred in Chesapeake Bay in the last 15 to 20 years (Orth and Moore 1983). Major changes in vegetation patterns began in 1972, the year of Tropical Storm Agnes, which lowered salinities for periods of up to 4 weeks and transported large quantities of suspended sediment into the estuarine system. The causes that have led to the Chesapeake Bay decline are not known but may be related to nutrient enrichment affecting the quantity and quality of light reaching the plant surface. Implications for species inhabiting grass beds have not been determined but could be considerable.

In Pinellas County and Hillsborough County, Florida, the two counties surrounding most of Tampa Bay, commercial spotted seatrout landings declined 65% from a 1951-55 average of 175 mt (386,000 lbs) to a 1976-80 average of 61 mt (135,000 lbs).² It has not been quantitatively demonstrated how these fish yields were affected by environmental changes, fishing pressure, or socio-economic changes. However, the environmental degradation in this area has been substantial (Taylor and Saloman 1968; Lewis 1977; Lewis and Phillips 1980). By comparing old and recent aerial photographs, Lewis and Phillips (1980) calculated that

²Pers. commun. Roy O. Williams, Florida Department of Natural Resources, St. Petersburg, Florida.

seagrass acreage on the Hillsborough County side of Tampa Bay declined 73% from 4,637 ha (11,458 acres) in 1948 to 1,251 ha (3,091 acres) in 1980. Concomitantly, commercial spotted seatrout landings in Hillsborough County declined 77% from a 93 mt (204,000 lbs) average during 1951-55 to 50 mt (110,000 lbs) during 1976-80.³ On the Pinellas County (St. Petersburg) side of the Bay, commercial spotted seatrout landings declined 59% from a 1951-56 average of 117 mt (257,000 lbs) to 48 mt (105,000 lbs) in 1976-80. No quantitative estimate of the habitat decline on the Pinellas County side of Tampa Bay is presently available. However, Taylor and Saloman (1968) estimated that in Boca Ciega Bay (a smaller bay off Tampa Bay within Pinellas County), 1,400 ha (3,500 acres) of bay bottom had been filled to create waterfront property for real estate development. This reduced the area of the bay by 20% and destroyed 1,133 mt of annual standing crop of seagrasses, resulting in an annual production loss of at least 73 mt (161,000 lbs) of fishery products.

In recent years the coastal states have enacted coastal zone management laws to regulate dredge and fill activities and shoreline development (Table 12-19). The Federal government also has some jurisdiction over the estuarine-marine habitat. The Office of Coastal Zone Management (OCZM) has authority through National Marine Sanctuaries, pursuant to Title III of the Marine Protection, Research, and Sanctuaries Act (MPRSA). The OCZM Estuarine Sanctuary program has designated Rookery Bay in Collier County, Florida, and the Apalachicola River and Bay in Franklin County, Florida, as estuarine sanctuaries. The OCZM also sets standards for approving and funding state coastal zone management programs. The Environmental Protection Agency may provide protection to fish communities through the granting of National Pollutant Discharge Elimination System (NPDES) permits for the discharge of pollutants into ocean waters, and the conditioning of those permits so as to protect valuable resources. The U. S. Army Corps of Engineers has jurisdiction over the disposal of dredged material, pursuant to both the Clean Water Act and the MPRSA. The Fish and Wildlife Service, under the Fish and Wildlife Coordination Act, reviews and comments on proposals for work and activities in or affecting navigable waters that are sanctioned, permitted, assisted, or conducted by Federal agencies. The review focuses mainly on potential damage to fish and wildlife, and their habitat.

The Environmental Assessment Branch of the NMFS is required to assess potential impacts on fishery resources of projects submitted to the Corps of Engineers for permits, and to recommend whether a project should be approved, denied, or modified. Fiscal year 1981 (October 1980 - September 1981) was the first year NMFS quantified the cumulative acreage of habitat involved in the Corps of Engineers permit program in the Southeast Region of the United States. NMFS made recommendations on 1,380 permit applications involving 7,272 ha (17,969 acres); 18% were proposed for dredging, 36% for filling, and 46% for impounding. NMFS did not object to alteration of 1,861 ha (4,598 acres), recommended against altering 5,411 ha (13,371 acres), and recommended that 1,345 ha (3,324 acres) either be restored or modified from upland habitat to

³Pers. commun. Roy O. Williams, Florida Department of Natural Resources, St. Petersburg, Florida.

Table 12-19. Summary of state habitat protection regulations, Rhode Island to Florida.

State	Administrative organization	Legislative authorization	Regulations
Rhode Island	Rhode Island Department of Environmental Management and Coastal Resources Management Council	Chapter 279, Public Laws of 1971, Sect. 1, Title 46, General Laws of Water and Navigation. Chapter 23 Coastal Resources Management Council.	Permits required for coastal zone development, aquaculture, dredge and fill operations.
Connecticut	Connecticut Department of Environmental Protection	"The Coastal Management Act" Section 22-a-90 to 22a-96	Permits required to dredge fill or construct structures in both fresh and salt water. Permit required to work in regulated wetland areas.
New York	Department of Environmental Conservation, Bureau of Tidal Wetlands	Environmental Conservation Law Article 25, Tidal Wetlands Act, Part 661. Land use regulations of tidal wetlands.	Regulates activities in and adjacent to tidal wetlands and requires permits for such activities.
New Jersey	Department of Environmental Protection, Division of Coastal Resources	Wetlands Act of 1970 NJSA 13:9A-1 et seq., Coastal Area Facilities Review Act NJSA 13:19-1 et seq., Waterfront Development Law, NJSA 12:5-3, Beaches and Harbors Bond Act of 1977 PL 77-208, Shore Protection Legislation NJSA 12:6A-1	Regulates activities in the coastal zone and requires permits for such activities.

Table 12-19. Continued

State	Administrative organization	Legislative authorization	Regulations
Delaware	Delaware Department of Natural Resources and Environmental Control, Division of Environmental Control, Wetlands Section	Sect. 1, Title 7, Delaware Code, Chapter 66. Wetlands.	Regulates use of wetlands and their upland border and provide penalties for violations.
Maryland	Maryland Department of Natural Resources, Tidewater Administration; Maryland Department of Health and Mental Hygiene, Office of Environmental Programs	Natural Resources Article, Code of Maryland	Regulates activities in tidal wetlands areas.
Virginia	Virginia Marine Resources Commission; County wetlands boards	The Wetlands Act Section 62.1-13.20, Code of Virginia	Regulates alterations to tidal marshes, sand and mud flats, subaqueous bottoms, and sand dunes.
North Carolina	North Carolina Department of Natural Resources and Community Development, Office of Coastal Management; Coastal Resources Commission; Coastal Resources Advisory Council	NC Dredge and Fill Law (GS 113-229), Coastal Area Management Act (CAMA) (GS 113A100)	Requires permits to dredge or fill in or about estuarine waters. Establishes areas of environmental concern. Permits required for coastal zone development.
	Division of Marine Fisheries	North Carolina Administrative Code, Chapter 3, Section .1400	Prohibits the use of bottom-disturbing gears and severely restricts or prohibits excavation and/or filling activities in nursery areas for young finfish and crustaceans.
South Carolina	South Carolina Coastal Zone Management Council	Coastal Zone Management and Planning Act	Directs permit activities in areas of wetlands,

Table 12-19. Continued

State	Administrative organization	Legislative authorization	Regulations
Georgia	Georgia Department of Natural Resources, Coastal Resources Division, Coastal Protection Section	Coastal Marshlands Protection Act of 1970 (Gs. L. 1970, p. 939, 1.)	Requires permits to dredge, fill, remove drain, or otherwise alter any marshlands.
		Shore Assistance Act of 1979 (Gs. L. 1979, 1.)	Requires permits for a structure, shoreline engineering activity, or land alteration in beaches, sand bars, and sand dunes in Georgia.
Florida	Florida Department of Natural Resources	Chapter 253, Florida Statutes	Regulates dredge, fill, and structures on state submerged lands (below mean high water). Provides for acquisition of conservation lands and tidally influenced areas.
		Chapter 258, F.S.	Establishes aquatic preserves and regulates activities within preserves.
	Florida Department of Environmental Regulation	Chapter 403, F.S.	Permitting of activities (including dredge and fill) which affect water quality.
	Florida Department of Community Affairs	Chapter 380, F.S.	Administer and set standards for "Development of Regional Impact". Protects regional or statewide resources from poorly conceived development activities.

mitigate the losses that were permitted. Thus, the NMFS efforts conserved 6,756 ha (16,695 acres) of habitat (Lindall and Thayer 1982). NMFS is also involved in the review of Congressionally authorized Federal projects. NMFS has adopted a new habitat conservation policy which will enhance its overall role in habitat conservation from a previously advisory role based primarily on the policies developed in response to the Fish and Wildlife Coordination Act and the National Environmental Policy Act.⁴ The new policy will: (1) ensure that habitat is fully considered in all of NMFS' programs and activities; (2) focus NMFS' habitat conservation activities on species for which the agency has management or protection responsibilities under the MFCMA, the Marine Mammal Protection Act, and the Endangered Species Act; (3) lay the foundation for management and research cooperation on habitat issues; and (4) strengthen NMFS' partnerships with the states and the regional FMCs on habitat issues.

12.7.3 Stocking

Uses of artificially propagated sciaenids as a management tool include: (1) description of early life history stages; (2) bioassay; and (3) introduction of tagged known-age stocks to determine growth, migratory patterns, and exploitation rate (Tatum 1981). Biologists with the Texas Parks and Wildlife Department successfully induced spotted seatrout to spawn and pictorially documented recently hatched and developing larvae (Colura 1974). Arnold et al. (1976) developed and described methods and techniques to maintain adult spotted seatrout in captivity, to induce them to spawn repeatedly, and to culture the young in order to have eggs, larvae, and juveniles of known history for experimental purposes. Larvae were reared to an age of 3 days with a 75-80% survival and to 30 days with a 30% survival. Cannibalism and lack of proper food appeared to be the major problems in the mass production of spotted seatrout. Growth of spotted seatrout larvae in the laboratory in relation to temperature, prey species and abundance, and stocking densities in the laboratory were reported by Taniguchi (1979, 1981) and Houde and Taniguchi (1982). Juvenile spotted seatrout have been stocked in ponds in Louisiana (Sackett et al. 1979) and in Texas (Colura et al. 1976).

12.8 Current Research

There is little ongoing research on spotted seatrout on the Atlantic coast. The Maryland Tidewater Administration, the Virginia Institute of Marine Science, and the North Carolina Division of Marine Fisheries (NCDMF) conduct juvenile fish surveys in the estuaries to monitor numerous species. The NCDMF also samples the commercial fisheries in order to monitor adult finfish stocks and will begin an adult estuarine fish survey in 1984. A tagging and biological study of spotted seatrout was recently completed by the Georgia Coastal Resources Division. The Florida Department of Natural Resources is examining habitat loss in three Florida estuaries and changes in the fisheries of those estuaries.

⁴Federal Register 48(228):53142-53148, November 25, 1983.

The National Park Service samples juvenile fishes in Everglades National Park and conducts a creel survey of the recreational fishery. The Fish and Wildlife Service is tagging spotted seatrout in Everglades National Park. NMFS Southeast Fisheries Center has initiated a habitat utilization study of seagrass beds in Everglades National Park. NMFS conducts an annual marine recreational fishery statistics survey. Commercial landings statistics are collected by state and Federal port agents.

12.9 Identification of Problems

The ISFMP Sciaenid Technical Committee has agreed that spotted seatrout research needs are: (1) yield modeling; (2) habitat requirements; (3) effects of environmental factors on stock density; and (4) delineation of Atlantic coast stocks. Improved catch and effort statistics for both the commercial and recreational fisheries are needed to measure stock density. The usefulness of controlling fishing mortality and minimum size need to be examined.

13.0 ACKNOWLEDGEMENTS

I wish to thank members of the Interstate Fisheries Management Program (ISFMP) Sciaenid Technical Committee for their major contributions to the format and contents of this report. The members of the ISFMP Sciaenid Board and the South Atlantic State Federal Fisheries Management Board provided guidance in the development of the plan's goal and objectives. Membership rosters are included as an appendix to this report (Section 15.0). I thank Michael Street, North Carolina Division of Marine Fisheries, for his advice and review of the manuscript. Funding for the preparation of this report was provided by the National Marine Fisheries Service (NMFS) Southeast Regional Office, NOAA, under State/Federal Cooperative Agreement No. SF-13. Funds for printing were provided by NMFS - Northeast Regional Offices, NOAA, to the ISFMP under Cooperative Agreement No. NO-80-FA-00017. I thank Diana Willis for typing this report.

14.0 REFERENCES

- Adams, C. A.
1972. Food habits of juvenile pinfish (Lagodon rhomboides), silver perch (Bairdiella chrysoura), and spotted seatrout (Cynoscion nebulosus) of the estuarine zone near Crystal River, Florida. M. S. Thesis, Univ. Florida, Gainesville, 147 p.
- Adkins, G. and M. J. Bourgeois.
1982. An evaluation of gill nets of various mesh sizes. La. Dep. Wildl. Fish., Seafood Div., Tech. Bull. No. 36, 59 p.
- Adkins, G., J. Tarver, P. Bowman and B. Savoie.
1979. A study of the commercial finfish in coastal Louisiana. La. Dep. Wildl. Fish., Seafood Div., Tech. Bull. No. 29, 87 p.
- Anderson, C. L. and R. H. McNutt.
1973. Costs and returns in commercial fishing: mullet fishing in Florida - a case study. Fla. Sea Grant Prog., Mar. Adv. Prog. Bull. SUSF-SG-73-002. State Univ. Syst., Gainesville, Fla.
- Anderson, W. W.
1968. Fishes taken during shrimp trawling along the South Atlantic coast of the United States, 1931-1935. U. S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 570, 60 p.
- Anderson, W. W. and J. W. Gehringer.
1965. Biological-statistical census of the species entering fisheries in the Cape Canaveral area. U. S. Fish. Wildl. Serv., Spec. Sci. Rep. Fish. 514, 79 p.
- Anonymous.
1979. Consumer survey of Florida seafood products. Fla. Dep. Nat. Resour., Bur. Market Exten. Serv., Tallahassee, Fla.
- Anonymous.
1980. Marine recreational fishery statistics survey, Atlantic and Gulf coasts, 1979. U. S. Natl. Mar. Fish. Serv. Curr. Fish. Stat. No. 8063, 139 p.
- Anonymous.
1983a. 1983 World record game fishes. Freshwater, Saltwater, and Fly Fishing. International Game Fish Assoc., Fort Lauderdale, Fla.
- Anonymous.
1983b. For inshore action, go for spotted seatrout. Coastlines Georgia 6(1):16-18.

Anonymous.

1984. Marine recreational fishery statistics survey, Atlantic and Gulf coasts, 1979 (revised)-1980. U. S. Natl. Mar. Fish. Serv. Curr. Fish. Stat. No. 8322, 239 p.

Arnold, C. R., J. L. Lasswell, W. H. Bailey, T. D. Williams, and W. A. Fable, Jr.

1976. Methods and techniques for spawning and rearing spotted seatrout in the laboratory. Proc. Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies 30:167-178.

Arnott, H. J., J. C. Nicol, and C. W. Querfeld.

1971. Reflecting speares in the eyes of weakfishes (Sciaenidae). Nature 233:130-132.

Bayley, P. B.

1977. A method for finding the limits of application of the von Bertalanffy growth model and statistical estimates of the parameters. J. Fish. Res. Board Can. 34:1079-1084.

Bearden, C. M.

1961. Common marine fishes of South Carolina. Bears Bluff Lab. Contrib. 34, 47 p.
1969. A report and recommendations on the saltwater sport fisheries of South Carolina. S. C. Wildl. Resour. Dep. unpub. ms, 96 p.

Bearden, C. M. and C. H. Farmer, III.

1972. Fishery resources of Port Royal Sound estuary. In Port Royal Sound Environmental Study, p. 203-212. S. C. Water Resour. Comm., 555 p.

Beaumariage, D. S.

1964. Returns from the 1963 Schlitz tagging program. Fla. Board Conserv., Mar. Res. Lab. Tech Ser. 43, 34 p.
1969. Returns from the 1965 Schlitz tagging program, including a cumulative analysis of previous results. Fla. Dep. Nat. Resour., Mar. Res. Lab. Tech Ser. No. 59, 38 p.

Beaumariage, D. S. and A. C. Wittich.

1966. Returns from the 1964 Schlitz tagging program. Fla. Board Conserv., Mar. Res. Lab. Tech. Ser. No. 47, 50 p.

Bigelow, H. B. and W. C. Schroeder.

1953. Fishes of the Gulf of Maine. U. S. Fish Wildl. Serv. Fish. Bull. 53:417-423.

- Breuer, J. P.
1973. A survey of the juvenile and adult food and game fish of the Laguna Madre. Tex. Parks Wildl. Dep., Coast. Fish. Prog. Rep. 1973:173-202.
- Brice, J. J., B. W. Evermann, B. A. Bean, and W. A. Wilcox.
1898. Report on the fisheries of Indian River, Florida. U. S. Comm. Fish Fish., Pt. XXII:223-262.
- Brown, N. J.
1981. Reproductive biology and recreational fishery for spotted seatrout, Cynoscion nebulosus, in the Chesapeake Bay area. M. A. Thesis, College of William and Mary, Williamsburg, 119 p.
- Brown, N. J., P. Thomas, and C. R. Arnold.
1983. Different life history strategies of the spotted seatrout, Cynoscion nebulosus, in a Texas and a Virginia estuary. (Abstract). Estuaries 6(3):260.
- Butler, P.A.
1969. The significance of DDT residues in estuarine fauna. In M.W. Miller and G. B. Berg (editors), Chemical Fallout, p. 205-220. Charles C. Thomas Publishers, Springfield, Ill.
- Butler, P. A., R. Childress and A. J. Wilson, Jr.
1970. The association of DDT residues with losses in marine productivity. FAO Technical Conference on Marine Pollution and its Effects on Living Resources and Fishing, 13 p.
- Cain, R. L. and J. M. Dean.
1976. Annual occurrence, abundance and diversity of fish in a South Carolina intertidal creek. Mar. Biol. 36:369-379.
- Carr, W. E. S. and C. A. Adams.
1973. Food habits of juvenile marine fishes occupying seagrass beds in the estuarine zone near Crystal River, Florida. Trans. Am. Fish. Soc. 102(3):511-540.
- Cato, J. C.
1981. Economic values and uses of the sciaenid fisheries. In Marine Recreational Fisheries 6. Proc. 6th Annu. Mar. Rec. Fish. Sym., p. 59-68. Sport Fishing Institute, Wash., D.C., 216 p.
- Chao, L. N.
1976. Aspects of systematics, morphology, life history, and feeding of western Atlantic Sciaenidae (Pisces: Perciformes). Ph. D. Diss., College of William and Mary, Williamsburg, 342 p.
1978. A basis for classifying western Atlantic Sciaenidae (Teleostei: Perciformes). U. S. Dep. Commer. NOAA Tech. Rep. NMFS Circ. 415, 64 p.

- Christmas, J. Y. and R. S. Waller.
1973. Cooperative Gulf of Mexico estuarine inventory and study, Mississippi. Phase IV: Biology. Section 5. Estuarine vertebrates, Mississippi. Miss. Mar. Conserv. Comm.:320-406.
- Clark, J. R.
1962. The 1960 salt-water angling survey. U. S. Bur. Sport Fish. Wildl. Circ. 153, 36 p.
- Colura, R.
1974. Induced spawning of the spotted seatrout Cynoscion nebulosus (Cuvier). Proc. Annu. Meet. World Mariculture Soc. 5:319-326.
- Colura, R. L., B. T. Hysmith, and R. E. Stevens.
1976. Fingerling production of striped bass (Morone saxatilis), spotted seatrout (Cynoscion nebulosus), and red drum (Sciaenops ocellatus), in saltwater ponds. Proc. Annu. Meet. World Mariculture Soc. 7:79-92.
- Condrey, R. E., G. Adkins and M. W. Wascom.
1984. A yield-per-recruit analysis of spotted seatrout. La. State Univ., Coastal Ecol. Fish. Inst., In-House CEFI Fish. Rep. No. 84-6, 17 p.
- Cupka, D. M.
1972. Winter trout. S. C. Wildl. 19:25-27.
- Dahlberg, M. D.
1972. An ecological study of Georgia coastal fishes. U. S. Natl. Mar. Fish. Serv. Fish. Bull. 70(2):323-353.
- Daniels, K. L.
1977. Description, comparison, and distribution of larvae of Cynoscion nebulosus and Cynoscion arenarius from the northern Gulf of Mexico. M.S. Thesis, La. State Univ., Baton Rouge, 48 p.
- Darnell, R. M.
1958. Food habits of fishes and larger invertebrates of Lake Pontchartrain, Louisiana, an estuarine community. Publ. Inst. Mar. Sci., Univ. Tex. 5:353-416.
- Davis, G. E.
1980. Changes in the Everglades National Park red drum and spotted seatrout fisheries 1958-1978: Fishing pressure, environmental stress, or natural cycles? Proc. Colloq. on the biology and management of red drum and seatrout. Gulf States Mar. Fish. Comm. Spec. Rep. No. 5:81-87.
1982. Fishery management conflicts in Everglades National Park. In Marine Recreational Fisheries 7. Proc. 7th Annu. Mar. Rec. Fish. Sym., p.65-75. Sport Fishing Institute, Wash., D. C., 183 p.

- Deuel, D. G.
1973. The 1970 salt-water angling survey. U. S. Natl. Mar. Fish. Serv. Curr. Fish. Stat. 6200, 54 p.
- Deuel, D. G. and J. Clark.
1968. The 1965 salt-water angling survey. U. S. Bur. Sport Fish. Wildl. Resour. Publ. 67, 51 p.
- DeVries, D. A.
1981. Description and catch composition of North Carolina's long haul seine fishery. Proc. Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies 34:234-274.
- Fable, W. A., Jr., T. D. Williams and C. R. Arnold.
1978. Description of reared eggs and young larvae of the spotted seatrout Cynoscion nebulosus. U. S. Natl. Mar. Fish. Serv. Fish. Bull. 76:65-71.
- Fontenot, B. J., Jr. and H. E. Rogillio.
1970. A study of estuarine sportfishes in the Biloxi Marsh Complex, Louisiana. La. Wildl. Fish. Comm. Fish Bull. No. 8, 172 p.
- Freeman, B. L. and L. A. Walford.
1974. Angler's guide to the United States Atlantic coast. Sect. IV. Delaware Bay to False Cape, Virginia. U. S. Govt. Print. Off., Wash., D. C., 17 p.
- 1976a. Angler's guide to the United States Atlantic coast. Sect. V. Chesapeake Bay. U. S. Govt. Print. Off., Wash., D. C., 17 p.
- 1976b. Angler's guide to the United States Atlantic coast. Sect. VI. False Cape, Virginia to Altamaha Sound, Georgia. U. S. Govt. Print. Off., Wash., D. C., 21 p.
- 1976c. Angler's guide to the United States Atlantic coast. Sect. VII. Altamaha Sound, Georgia to Fort Pierce Inlet, Florida. U. S. Govt. Print. Off., Wash., D. C., 17 p.
- 1976d. Angler's guide to the United States Atlantic coast. Section VII. St. Lucie Inlet, Florida to the Dry Tortugas. U. S. Govt. Print. Off., Wash., D. C., 21 p.
- Futch, C. R.
1970. The spotted seatrout, Cynoscion nebulosus (Cuvier). Fla. Dep. Nat. Resour. Salt Water Fish. Leaflet. 11, 11 p.
- Gadbois, D. F. and R. S. Maney.
1983. Survey of polychlorinated biphenyls in selected finfish species from United States coastal waters. U. S. Natl. Mar. Fish. Serv. Fish. Bull. 81(2):389-396.

- Gilmore, R. G., Jr.
1977. Fishes of the Indian River lagoon and adjacent waters, Florida. Bull. Fla. State Mus. Biol. Sci. 22(3):101-148.
- Gilmore, R. G., G. R. Kulezycki, P. A. Hastings, and W. C. Magley.
1976. Chapt. 11. Studies of fishes of the Indian River Lagoon and vicinity. p. 133-147. In Indian River Coastal Zone Study, 1975-1976 Annu. Rep. Vol. 1, p. 133-147. Harbor Branch Consortium, 187 p.
- Ginsburg, I.
1929. Review of the weakfishes (Cynoscion) of the Atlantic and Gulf Coasts of the United States with a description of a new species. Bull. U. S. Bur. Fish. 45:71-85.
- Goode, G. B.
1884. The fisheries and fishery industries of the United States. Section I. Natural History of Useful Aquatic Animals. Text, p. 365-367. Govt. Print. Off., Wash.

1887. The fisheries and fishery industries of the United States. II. A geographical review of the fisheries industries and fishing communities for the year 1880. Rep. U. S. Comm. Fish, Fish. Sect. II:526-531.
- Greenwood, P. D. Rosen, W. Weitzman and G. Myers.
1966. Phyletic studies of teleostean fishes, with a provisional classification of living forms. Bull. Am. Mus. Nat. Hist. 131:341-455.
- Guest, W. C. and G. Gunter.
1958. The seatrout or weakfishes (Genus Cynoscion) of the Gulf of Mexico. Gulf States Mar. Fish. Comm., Tech. Summ. No. 1, 40 p.
- Gunter, G.
1938. Seasonal variations in abundance of certain estuarine and marine fishes in Louisiana, with particular reference to life histories. Ecol. Monogr. 8(3):314-346.

1941. Death of fishes due to the cold on the Texas coast. Ecology 22(2):203-308.

1945. Studies on marine fishes in Texas. Publ. Inst. Mar. Sci., Univ. Tex. 1(1):1-190.
- Gunter, G. and H. H. Hildebrand.
1951. Destruction of fishes and other organisms on the south Texas coast by the cold wave of January 28-February 3, 1951. Ecology 32(4):731-735.
- Gusey, W. F.
1978. The fish and wildlife resources of the Middle Atlantic Bight. Shell Oil Company, Houston, Tex.

1981. The fish and wildlife resources of the South Atlantic coast. Shell Oil Company, Houston, Tex.
- Hall, R. A., E. G. Zook and G. M. Meaburn.
1978. National Marine Fisheries Service survey of trace elements in the fishery resource. NOAA Tech. Rep. NMFS SSRF-721, 313 p.
- Hammond, D. L. and D. M. Cupka.
1977. An economic and biological evaluation of the South Carolina pier fishery. S. C. Mar. Resour. Cent. Tech. Rep. No. 20, 14 p.
- Harrington, R. A., G. C. Matlock and J. E. Weaver.
1979. Standard-total length, total length-whole weight, and dressed-whole weight relationships for selected species from Texas bays. Tex. Parks Wildl. Dep. Tech. Ser. 26, 6 p.
- Heffernan, T. L. and R. J. Kemp.
1982. The conflicts and controversies surrounding red drum and spotted seatrout. In Marine Recreational Fisheries 7. Proc. 7th Annu. Mar. Rec. Fish. Sym., p.57-64. Sport Fishing Institute, Wash., D.C., 183 p.
- Hein, S. H. and J. A. Shepard.
1979a. Spawning of spotted seatrout in a Louisiana estuarine ecosystem. Proc. Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies 33:451-465.
1979b. Size disparity between "Inside" and "Outside" Spotted Seatrout (Cynoscion nebulosus) during a three month study in south-central Louisiana. La. Dep. Wildl. Fish., Contrib. Mar. Res. Lab., Tech. Bull. No. 28:16-28.
1980. Size of spotted seatrout (Cynoscion nebulosus) captured in three and one-quarter inch stretched monofilament gill nets in South Louisiana. La. Dep. Wildl. Fish., Contrib. Mar. Res. Lab. - 1978, Tech. Bull. No. 31:49-54.
- Hein, S., C. Dungas and J. Shepard.
1980. Total length-standard length and length-weight regressions for spotted seatrout, Cynoscion nebulosus; red drum, Sciaenops ocellata; and black drum, Pogonias cromis, in South-central Louisiana. La. Dep. Wildl. Fish., Contrib. Mar. Res. Lab. - 1978, Tech. Bull. No. 31:41-48.
- Hicks, D. B.
1972. Seasonal distribution and relative abundance of fishes in the channel reaches and shore areas. In Port Royal Sound Environmental Study, p. 193-201. S. C. Water Resour. Comm., 555 p.

- Higgins, E. and J. C. Pearson.
1928. Examination of the summer fisheries of Pamlico and Core Sounds, N. C., with special reference to the destruction of undersized fish and the protection of the gray trout, *Cynoscion regalis* (Bloch and Schneider). Rep. U.S. Comm. Fish., 1927 Appendix II:29-65.
- Higman, J. B.
1967. Relationship between catch rates of sport fish and environmental conditions in Everglades National Park, Florida. Proc. 19th Annu. Sess. Gulf Caribb. Fish. Inst. 19:129-140.
- Hildebrand, S. F. and L. E. Cable.
1934. Reproduction and development of whittings or kingfishes, drums, spot, croaker, and weakfishes or seatrouts, family Sciaenidae, of the Atlantic coast of the United States. Bull. U. S. Bur. Fish. 48:41-117.
- Hildebrand, S. F. and W. C. Schroeder.
1928. Fishes of Chesapeake Bay. Bull. U. S. Bur. Fish. 43(1):296-299.
- Hoese, H. D.
1973. A trawl study of nearshore fishes and invertebrates of the Georgia coast. Contrib. Mar. Sci. 17:63-98.
- Hoese, H. D. and R. H. Moore.
1977. Fishes of the Gulf of Mexico, Texas, Louisiana, and adjacent waters. Tex. A & M Univ. Press, 327 p.
- Holt, G. J., S. A. Holt, and C. R. Arnold.
1983. Spawning synchrony in sciaenid fishes. (Abstract). Estuaries 6(3):261.
- Houde, E. D. and A. K. Taniguchi.
1982. Marine fish larvae growth and survival. U. S. Environ. Prot. Agency, Environ. Res. Lab., Narragansett, R. I., Res. Devel. EPA-600/53-81-052.
- Houde, E. D., J. C. Leak, C. E. Dowd, S. A. Berkley, and W. J. Richards.
1979. Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. Rep. to Bur. Land Manag., Contract No. AA550-CT7-28:35.
- Idyll, C. P. and W. E. Fahy.
1970. Spotted seatrout. . . shallow-water sport fish. Mar. Resour. Atl. Coast, Atlantic States Mar. Fish. Comm. Leaflet 13, 4 p.
- Ingle, R. M., R. F. Hutton and R. W. Topp.
1962. Results of the tagging of salt water fishes in Florida. Fla. Board Conserv. Mar. Res. Lab. Tech. Ser. 38:1-57.

- Iversen, E. S. and A. W. Moffett.
1962. Estimates of abundance and mortality of a spotted seatrout population. *Trans. Am. Fish. Soc.* 91(4):395-398.
- Iversen, E. S., and D. C. Tabb.
1962. Subpopulations based on growth and tagging studies of spotted seatrout, Cynoscion nebulosus, in Florida. *Copeia* (1962):544-548.
- Jannke, T. E.
1971. Abundance of young sciaenid fishes in Everglades National Park, Florida in relation to season and other variables. *Univ. Miami Sea Grant Tech. Bull.* 11, 128 p.
- Johnson, A. G., T. D. Williams, and C. R. Arnold.
1977. Chlorine-induced mortality of eggs and larvae of spotted seatrout (Cynoscion nebulosus). *Trans. Am. Fish. Soc.* 106(5):466-469.
1979. Larval spotted seatrout (Cynoscion nebulosus) a bioassay subject for the marine subtropics. *Contrib. Mar. Sci.*, 22:57-62.
- Johnson, G. D.
1978. Development of fishes of the Mid-Atlantic Bight. In *An atlas of egg, larval and juvenile stages. Vol. IV. Carangidae through Ehippidae.* 180:186. U. S. Fish. Wildl. Serv. Biol. Serv. Program FWS/OBS-78/12.
- Jones, R. S., R. G. Gilmore, Jr., G. R. Kulczycki, W. C. Magley and B. Graunke.
1975. Chapter 4. Studies of the Fishes of the Indian River coastal zone. In *Indian River Coastal Zone Study, 1974-75 Annu. Rep. Vol. I.* p. 57-88. Harbor Branch Consortium, 180 p.
- Jordan, D. S. and B. W. Evermann.
1896. The fishes of North and Middle America. *U. S. Nat. Mus. Bull.* 47. 1240 p.
- Jorgenson, S. C. and G. L. Miller.
1968. Length relations of some marine fishes from coastal Georgia. *U. S. Fish Wildl. Serv. Spec. Sci. Rep. Fish. No.* 575, 16 p.
- Juneau, C. L.
1975. An inventory and study of the Vermillion Bay - Atchafalaya Bay complex. *La. Wildl. Fish. Comm. Tech. Bull.* No. 13, 153 p.
- Keiser, R. K., Jr.
1976. Species composition, magnitude and utilization of the incidental catch of the South Carolina shrimp fishery. *S. C. Mar. Resour. Cent. Tech. Rep. No.* 16, 55 p.

- Kemp, R. J.
1949. Report on stomach analyses from June 1, 1949. Tex. Game, Fish, Oyst. Comm., Mar. Lab. Annu. Rep. 1948-49:101-127.
- King, B. D., III.
1971. Study of migratory patterns of fish and shellfish through a natural pass. Tex. Parks Wildl. Dep., Tech. Ser. 9. 54 p.
- Klima, E. F. and D. C. Tabb.
1959. A contribution to the biology of the spotted weakfish, Cynoscion nebulosus (Cuvier), from northwest Florida, with a description of the fishery. Fla. Board Conserv. Mar. Res. Lab. Tech. Ser. 30, 25 p.
- Knapp, F. T.
1949. Menhaden utilization in relation to the conservation of food and game fishes on the Texas Gulf coast. Trans. Am. Fish. Soc. 79:137-144.
- Knowlton, C. J.
1972. Fishes taken during commercial shrimp fishing in Georgia's close inshore ocean waters. Ga. Game Fish Comm., Coast. Fish. Off., Contrib. Ser. No. 21, 42 p.
- Lascara, J.
1981. Fish predator-prey interactions in areas of eelgrass (Zostera marina). M.A. Thesis, College of William and Mary, Williamsburg, 81 p.
- Lewis, R. R.
1977. Impacts of dredging in the Tampa Bay estuary, 1976-1976. In E.L. Pruitt (editor), Proc. 2nd Annu. Conf. Coastal Soc., Nov. 17-20, 1976; New Orleans, La., p. 31-55. The Coastal Society, Arlington, Va.
- Lewis, R. R. and R. C. Phillips.
1980. Seagrass Mapping Project, Hillsborough County, Florida. Final Report (mimeo). Mangrove Systems, Inc., Tampa, 15 p.
- Lindall, W. N., Jr. and G. W. Thayer.
1982. Quantification of National Marine Fisheries Service habitat conservation efforts in the Southeast Region of the United States. U. S. Natl. Mar. Fish. Serv. Mar. Fish. Rev. 44 (12):18-22.
- Linton, E.
1905. Parasites of fishes of Beaufort, North Carolina. Bull. U. S. Bur. Fish. 24:321-428.
- Lippson, A. J. and R. L. Moran.
1974. Manual for identification of early development stages of fishes of the Potomac River estuary. Md. Dep. Nat. Resour. PPSP-MP-13, 282 p.

- Loman, M.
1978. Other finfish. In Fisheries assessment and monitoring Mississippi, p. 143-147. Gulf Coast Research Lab. Complet. Rep. P.L. 88-309, 2-215-R.
- Lorio, W. J. and H. E. Schafer.
1966. A food habit study of the spotted seatrout, Cynoscion nebulosus, in the Biloxi Marsh area, Louisiana. Proc. Annu. Conf. Southeast. Assoc. Game Fish Comm. 19:289-296.
- Lorio, W. J. and W. S. Perret.
1980. Biology and ecology of the spotted seatrout (Cynoscion nebulosus Cuvier). Proc. Colloq. on the biology and management of red drum and seatrout, Gulf States Mar. Fish. Comm. Spec. Rep. No. 5:7-13.
- Lunz, G. R. and F. J. Schwartz.
1969. Analysis of eighteen year trawl captures of seatrout (Cynoscion sp.: Sciaenidae) from South Carolina. Contrib. Bears Bluff Lab. No. 53, 29 p.
- Mahood, R. K.
1974. Seatrout of the genus Cynoscion in coastal waters of Georgia. Ga. Dep. Nat. Resour. Contrib. Ser. 26, 35 p.
1975. Spotted seatrout in coastal waters of Georgia. Proc. Annu. Conf. Southeast. Assoc. Game Fish. Comm. 29:195-207.
- Mather, F. J., III.
1952. Sport fishes of the vicinity of the Gulf of Honduras, certain Caribbean islands, and Carmen, Mexico. Proc. Gulf. Caribb. Fish. Inst. 4:118-129.
- Matlock, G. C.
1982. The conflict between user groups of red drum and spotted seatrout in Texas. In Marine Recreational Fisheries Proc. 7th Annu. Mar. Rec. Fish. Sym., p. 101-108. Sport Fishing Institute, Wash., D.C., 183 p.
- Matlock, G. C., J. E. Weaver, and A. W. Green.
1979. Trends in spotted seatrout and red drum abundance in Texas coastal waters influenced by commercial netting activities. Proc. Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies 31:477-483.
- Matlock, G. C. and M. A. Garcia.
1983. Stomach contents of selected fishes from Texas bays. Contrib. Mar. Sci. 26:95-110.
- Mercer, L. P.
1984. A biological and fisheries profile of spotted seatrout, Cynoscion nebulosus. N. C. Div. Mar. Fish. Spec. Sci. Rep. No. 40, 87 p.

Merriner, J. V.

1980. History and management of the spotted seatrout fishery. Proc. Colloq. on the biology and management of red drum and seatrout. Gulf States Mar. Fish. Comm. Spec. Rep. No. 5:55-61.

Miles, D. W.

1949. A study of the food habits of the fishes of the Aransas Bay area. Tex. Game Fish Oyster Comm. Mar. Lab. Annu. Rep., 1948-49:129-169.

1950. The life histories of spotted seatrout, Cynoscion nebulosus, and the redfish, Sciaenops ocellatus. Tex. Game Fish. Comm., Mar. Lab. Annu. Rep., 1949-1950, 30 p.

1951. The life histories of the seatrout, Cynoscion nebulosus, and the redfish, Sciaenops ocellatus: sexual development. Tex. Game Fish. Comm., Mar. Lab. Annu. Rep., 1950-1951, 11 p.

Miller, G. L. and S. C. Jorgenson.

1969. Seasonal abundance and length frequency distribution of some marine fishes in coastal Georgia. U. S. Fish Wildl. Serv. Data Rep. 35, 102 p.

1973. Meristic characters of some marine fishes of the western Atlantic Ocean. U. S. Nat. Mar. Fish. Serv. Fish. Bull. 71:301-317.

Miller, J. M. and M. L. Dunn.

1980. Feeding strategies and patterns of movement in juvenile estuarine fishes. In V.S. Kennedy (editor), Estuarine Perspectives, p. 437-448. Acad. Press, N.Y., 533 p.

Moffett, A. W.

1961. Movements and growth of spotted seatrout, Cynoscion nebulosus (Cuvier), in west Florida. Fla. Board Conserv., Mar. Res. Lab., Tech. Ser. 36, 35 p.

Mohsin, A. K. M.

1973. Comparative osteology of the weakfishes (Cynoscion) of the Atlantic and Gulf coasts of the United States. Ph. D. Diss., Texas A & M Univ., College Station, 148 p.

Mok, H. K. and R. G. Gilmore.

1983. Analysis of sound production in estuarine aggregations of Pogonias cromis, Bairdiella chrysoura, and Cynoscion nebulosus (Sciaenidae). Bull. Inst. Zool., Acad. Sin. (Taipei) 22(2):157-186.

Moody, W. D.

1950. A study of the natural history of spotted trout, Cynoscion nebulosus, in the Cedar Key, Florida area. Q. J. Fla. Acad. Sci. 12:147-171.

- Moore, C. J.
1980. South Carolina's noncommercial gill-net fishery. Trans. Am. Fish. Soc. 109:577-580.
- Moore, R. H.
1976. Observations on fishes killed by cold at Port Aransas, Texas, 11-12 January 1973. Southwest Nat. 20:461-466.
- Music, J. L., Jr.
1981. Seasonal movement and migration of spotted seatrout (Cynoscion nebulosus) (Abstract). Estuaries 4(3):280.
- Music, J. L., Jr. and J. M. Pafford.
1984. Population dynamics and life history aspects of major marine sportfishes in Georgia's coastal waters. Ga. Dep. Nat. Resour., Coastal Resour. Div., Coastal Fish. Sect., Final Rep. Study VI. Fed. Aid Proj. F-31.
- Odum, W. E.
1971. Pathways and energy flow in a south Florida estuary. Sea Grant Tech. Bull. No. 7, 62 p.
- Odum, W. E. and E. J. Heald.
1972. Trophic analysis of an estuarine mangrove community. Bull. Mar. Sci. 22(3):671-738.
- Orth, R. J. and K. A. Moore.
1983. Chesapeake Bay: An unprecedented decline in submerged aquatic vegetation. Science 222:51-53.
- Orth, R. J. and K. L. Heck, Jr.
1980. Structural components of eelgrass (Zostera marina) meadows in the lower Chesapeake Bay - fishes. Estuaries 3(4):278-288.
- Osborne, K.
1981. Saltwater sport fishing and boating in North Carolina. Alexandria Drafting Co., Alexandria, Va., 52 p.
- Overstreet, R. M.
1983. Aspects of the biology of the spotted seatrout, Cynoscion nebulosus, in Mississippi. Gulf Res. Rep., Supp. 1:1-43.
- Overstreet, R. M. and R. W. Heard.
1982. Food contents of six commercial fishes from Mississippi Sound. Gulf Res. Rep. 7(2):137-149.
- Pearson, J. C.
1929. Natural history and conservation of the redfish and other commercial sciaenids on the Texas coast. Bull. U. S. Bur. Fish. 64:178-194.

- Perret, W. S.
1971. Phase IV, Biology. In Cooperative Gulf of Mexico estuarine inventory and study, Louisiana. p. 41-105. La. Wildl. Fish. Comm. Proj. 2-22R.
- Perret, W. S., J. E. Weaver, R. L. Williams, P. L. Johansen, T. D. McIlwain, R. C. Raulerson and W. M. Tatum.
1980. Fishery profiles of red drum and spotted seatrout. Gulf States Mar. Fish. Comm. No. 6, 60 p.
- Powles, H. and B. W. Stender.
1978. Taxonomic data on the early life history stages of Sciaenidae of the South Atlantic Bight of the United States. S. C. Mar. Resour. Cent. Tech. Rep. No. 31, 64 p.
- Purvis, C.
1976. Nursery area survey of northern Pamlico Sound and tributaries. N. C. Div. Mar. Fish. Complet. Rep., Proj. No. 2-230-R, 62 p.
- Randolph, J.
1983. Featured Fish: the spotted seatrout. The Virginia-North Carolina Fishing Report, Oct. 4:4.
- Reid, S. K., Jr.
1954. An ecological study of the Gulf of Mexico fishes, in the vicinity of Cedar Key, Florida. Bull. Mar. Sci. Gulf Caribb. 4(1):1-94.
- Renfro, W. C.
1963. Gas bubble mortality of fishes in Galveston Bay, Texas. Trans. Am. Fish. Soc. 92:320-322.
- Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooks, E. A. Lachner, R. N. Lea, and W. B. Scott.
1980. A list of common and scientific names of fishes from the United States and Canada. Am. Fish. Soc. Spec. Pub. No. 12, 174 p.
- Robison, D. E.
(in press). Variability in the vertical distribution of ichthyoplankton in lower Tampa Bay, Florida. Proc. Bay Area Scientific Information Symposium, Tampa Bay, Fla.
- Roelofs, E. W.
1951. The Edible Finfishes of North Carolina. In H. F. Taylor, Survey of Marine Fisheries of North Carolina, p. 127-128. Univ. N. C. Press, Chapel Hill, 555 p.
- Rogillio, H. E.
1975. An estuarine sportfish study in southeastern Louisiana. La. Wildl. Fish. Comm., Fish. Bull. No. 14, 71 p.

- Rosen, A. and R. Ellis.
1958. Summary of Florida commercial marine fish landings. Fla. Board Conserv., Univ. Miami Mar. Lab., Mar. Fish. Res., 58+2, 65 p.
- Rutherford, E. S.
1982. Age, growth and mortality of spotted seatrout, Cynoscion nebulosus, in Everglades National Park, Florida. M.S. Thesis, Univ. Miami, Coral Gables, 65 p.
- Rutherford, E. S., E. Thue and D. Buker.
1982. Population characteristics, food habits and spawning activity of spotted seatrout, Cynoscion nebulosus, in Everglades National Park, Florida. U. S. Natl. Park Serv., S. Fla. Res. Cent. Rep. T-668, 48 p.
- Sabins, D. S. and F. M. Truesdale.
1974. Diel and seasonal occurrence of immature fishes in a Louisiana tidal pass. Proc. Annu. Conf. Southeast. Assoc. Game Fish. Comm. 28:161-171.
- Sackett, S., S. H. Hein, and R. L. Hooton.
1979. Probable natural spawn and growth of juvenile spotted seatrout (Cynoscion nebulosus) in a 1/4-acre pond. La. Dep. Wildl. Fish. Contrib. Mar. Res. Lab. 1977, Tech. Bull. No. 28:54.
- Seagle, J. H.
1969. Food habits of spotted seatrout Cynoscion nebulosus (Cuvier) frequenting turtle grass (Thalassia testudinum, Konig) beds in Redfish Bay, Texas. Taius 2(1):58-63.
- Setzler, E. M.
1977. A quantitative study of the movement of larval and juvenile Sciaenidae and Engraulidae into the estuarine nursery grounds of Doboy Sound, Sapelo Island, Georgia. Ph. D. Diss., University of Georgia, Athens, 121 p.
- Shealy, M. H., J. V. Miglarese and E. B. Joseph.
1974. Bottom fishes of South Carolina estuaries, relative abundance, seasonal distribution and length-frequency relationships. S. C. Mar. Resour. Cent. Tech. Rep. No. 6, 189 p.
- Shiino, S. M.
1976. List of common names of fishes of the world, those prevailing among English-speaking nations. Shima Marineland, Kashikojima, Shima, Mie, Japan, 262 p.
- Simmons, E. G.
1951. Fish trap investigation. Tex. Game Fish Comm., Mar. Lab. Annu. Rep. 1950-1951. 23 p.

1957. Ecological survey of the upper Laguna Madre of Texas. Publ. Inst. Mar. Sci., Univ. Tex. 4:156-200.
- Smith, H. M.
1907. The Fishes of North Carolina. N. C. Geol. Econ. Survey 2, 449 p.
- Smith, J. W. and C. J. Moore.
1979. Coastal finfish management planning profile. S. C. Wildl. Mar. Resour. Dep., Off. Conserv., Manage., Market. unpub. ms, 140 p.
- Spitsbergen, D. L. and M. Wolff.
1974. Survey of nursery areas in western Pamlico Sound, North Carolina. N. C. Div. Mar. Fish. Complet. Rep. Proj. No. 2-175-R, 80 p.
- Springer, V. G. and K. D. Woodburn.
1960. An ecological study of the fishes of the Tampa Bay area. Fla. Board Conserv. Mar. Lab. Prof. Pap. Ser. 1:1-104.
- Stewart, K. W.
1961. Contributions to the biology of the spotted seatrout (Cynoscion nebulosus) in the Everglades National Park, Florida. M.S. Thesis, Univ. of Miami, Coral Gables, 103 p.
- Storey, M. and E. W. Gudger.
1936. Mortality of fishes due to cold at Sanibel Island, Florida, 1886-1936. Ecology 17(4):640-648.
- Sundararaj, B. I. and R. D. Suttkus.
1962. Fecundity of the spotted seatrout, Cynoscion nebulosus (Cuvier), from Lake Borgne area, Louisiana. Trans. Am. Fish. Soc. 9:84-84.
- Tabb, D. C.
1958. Differences in the estuarine ecology of Florida waters and their effect on populations of the spotted weakfish, Cynoscion nebulosus (Cuvier and Valenciennes). Proc. North Am. Wildl. Conf. 23:329-401.

1960. The spotted seatrout fishery of the Indian River area, Florida. Fla. Board Conserv. Tech. Ser. 33, 20 p.

1961. A contribution to the biology of the spotted seatrout, Cynoscion nebulosus (Cuvier), of east-central Florida. Fla. Board Conserv. Mar. Res. Lab. Tech. Ser. 35, 24 p.

1966. The estuary as a habitat for spotted seatrout, Cynoscion nebulosus. Am. Fish. Soc. Spec. Publ. 3:58-67.

Tabb, D. C. and R. B. Manning.

1961. A checklist of the flora and fauna of northern Florida Bay and adjacent brackish waters of the Florida mainland collected during the period July, 1957 through September, 1960. Bull. Mar. Sci. Gulf Caribb. 11(4):552-649.

Tagatz, M. E.

1967. Fishes of the St. Johns River, Florida. Q. J. Fla. Acad. Sci. 30(1):25-50.

Taniguchi, A. K.

1979. Survival and growth of larval spotted seatrout (Cynoscion nebulosus) larvae in relation to temperature, prey abundance and stocking densities. ICES/ELH Symp., 28 p.
1980. Effects of salinity, temperature and food abundance upon survival of spotted seatrout eggs and larvae. Proc. Colloq. on the biology and management of red drum and seatrout. (Abstract). Gulf State Mar. Fish. Comm. Spec. Rep. No. 5:16.
1981. Survival and growth of larval spotted seatrout (Cynoscion nebulosus) in relation to temperature, prey abundance and stocking densities. Rapp. P.-V. Reun. Cons. Int. Explor. Mer. 178:507-508.

Tatum, W. M.

1980. Spotted seatrout (Cynoscion nebulosus) age and growth: data from annual fishing tournaments in coastal Alabama, 1964-1977. Proc. Colloq. on the biology and management of red drum and seatrout. Gulf States Mar. Fish. Comm. Spec. Rep. No. 5:89-92.
1981. Utilizing artificially propagated sciaenids as a management tool. In Marine Recreational Fisheries 6, Proc. 6th Annu. Mar. Rec. Fish. Sym., p. 197-203. Sport Fishing Institute, Wash., D.C., 212 p.

Taylor, J. L. and C. H. Saloman.

1968. Some effects of hydraulic dredging and coastal development in Boca Ciega Bay, Florida. U. S. Natl. Mar. Fish. Serv. Fish. Bull. 67(2):213-241.

Thayer, G. W., H. H. Stuart, W. J. Kenworthy, J. F. Ustach, and A. B. Hall.

1978. Habitat values of salt marshes, mangroves, and seagrasses for aquatic organisms. In P.E. Greeson, J.R. Clark and J.E. Clark (editors), Wetland functions and values: the state of our understanding. Proc. Natl. Sym. Wetlands, p. 235-247. Am. Water Resour. Assoc.

- Topp, R.
1963. The tagging of fishes in Florida 1962 program. Fla. Dep. Nat. Resour. Mar. Res. Lab. Prof. Pap. Ser. 5, 76 p.
- Turner, W. R. and G. N. Johnson.
1972. Standing crops of aquatic organisms in five South Carolina tidal streams. In Port Royal Sound Environmental Study, p. 179-192. S. C. Water Resour. Comm.
- Vetter, R. D.
1977. Respiratory metabolism of, and niche separation between, two co-occurring congeneric species, Cynoscion nebulosus and Cynoscion arenarius, in a south Texas estuary. M.A. Thesis, Univ. Texas, Austin, 113 p.

1982. Seasonal metabolic compensation in sympatric seatrout: adaptation to the estuary. Trans. Am. Fish. Soc. 111:193-198.
- Wade, C. W.
(in press). Age and growth of spotted seatrout and red snapper in Alabama. Proc. Annu. Conf. Southeast. Assoc. Fish. Wildl. Agencies.
- Wakeman, J. M. and D. E. Wohlschlag.
1977. Salinity stress and swimming performance of spotted seatrout. Proc. Annu. Conf. Southeast. Assoc. Fish. Wildl. Agencies 31:357-361.
- Waller, R. S. and F. C. Sutter.
1982. Chapt. V. Section 1. Other. Fishery monitoring and assessment completion report. Gulf Coast Res. Lab. Proj. No. 2-296-R.
- Weaver, J. E.
1977. Recreational harvest of fishes in Texas bays. Tex. Parks Wildl. Dep., Coast. Fish. Branch. unpub. rep. presented to Am. Fish. Soc. Meet., Vancouver, B. C.
- Weinstein, M. P.
1981. Biology of adult sciaenids. In Marine Recreational Fisheries 6. 6th Annu. Mar. Rec. Fish. Sym., p. 125-138. Sport Fishing Institute, Wash., D.C., 212 p.
- Weinstein, M. P. and R. W. Yerger.
1976a. Protein taxonomy of the Gulf of Mexico and Atlantic Ocean seatrouts, genus Cynoscion. U. S. Natl. Mar. Fish. Serv., Fish. Bull., 74:599-607.

1976b. Electrophoretic investigation of subpopulations of the spotted seatrout, Cynoscion nebulosus (Cuvier), in the Gulf of Mexico and Atlantic coast of Florida. Comp. Biochem. Physiol. 54B:97-102.

- Welsh, W. W. and C. M. Breder, Jr.
1923. Contributions to life histories of Sciaenidae of the eastern United States coast. Bull. U. S. Bur. Fish. 39:141-201.
- Williams, A. B. and E. E. Deubler, Jr.
1968. Studies on macroplanktonic crustaceans and ichthyoplankton of the Pamlico Sound complex. N. C. Dep. Conserv. Sports Fish. Spec. Sci. Rep. No. 13, 103 p.
- Williams, J. B., H. J. Speir, S. Early, and T. P. Smith.
1982. 1979 Maryland saltwater sport fishing survey. Md. Dep. Nat. Resour. Tidewater Admin. TA-CRD-82-1, 100 p.
- Williams, J. B., T. P. Smith, H. J. Speir, and S. Early.
1983. 1980 Maryland saltwater sport fishing survey. Md. Dep. Nat. Resour. Tidewater Admin. TA-CRD-83-1, 124 p.
- Wohlschlag, D. E. and J. M. Wakeman.
1978. Salinity stresses, metabolic responses and distribution of the coastal spotted seatrout, Cynoscion nebulosus. Contrib. Mar. Sci. 21:171-185.
- Wolff, M.
1972. A study of the North Carolina scrap fishery. N. C. Div. Commer. Sport Fish. Spec. Sci. Rep. 20, 29 p.
1976. Nursery area survey of the Outer Banks region. N. C. Div. Mar. Fish. Complet. Rep. Proj. No. 2-222-R, 47 p.
- Yanez-Arancibia, A., F. A. Linares, and J. W. Day, Jr.
1980. Fish community structure and function in Terminos Lagoon, a tropical estuary in the southern Gulf of Mexico. In V. S. Kennedy (ed.) Estuarine Perspectives, p. 465-482. Acad. Press, N.Y., 533 p.
- Zieman, J. C.
1982. The ecology of the seagrasses of south Florida: a community profile. U. S. Fish Wildl. Serv. Biol. Serv. Prog. FWS/OBS - 82/25, 123 p.

15.0 APPENDIX

15.1. Listing of members of Sciaenid Technical Committee, Sciaenid Board,
and South Atlantic State/Federal Fisheries Management Board.Sciaenid Technical Committee

Chris Ordzie
RI Dept. Env. Manage.
Coastal Fish. Lab.
Succotash, RI
Wakefield, RI 02879
(401) 783-2304

George Maltezos
CT Dep. Env. Prot.
Marine Fish. Office
P.O. Box 248
Waterford, CT 06385
(203) 443-0166

Robert Sampson
CT Dep. Env. Prot.
Marine Fish. Office
P.O. Box 248
Waterford, CT 06385
(203) 443-0166

John Poole
NYS Dep. Env. Cons.
Bldg. 40, SUNY
Stony Brook, NY 11794
(516) 751-7900

Alice Weber
NYS Dep. Env. Cons.
Bldg. 40, SUNY
Stony Brook, NY 11794
(516) 751-7900

Paul Hamer
NJ Div. Fish, Game
& Wildlife
Mar. Fish. Lab.
Nacote Creek
Absecon, NJ 08201
(609) 441-3292

Richard Seagraves
DE Div. Fish. &
Wildlife
Edward Tatnal Bldg.
P.O. Box 1401
Dover, DE 19901
(301) 736-4782

Edward Houde
Univ. Maryland
Chesapeake Biological
Lab
P.O. Box 38
Solomons, MD 20688
(301) 326-4281

Charles Frisbie
MD Dept. Nat. Resources
Tidewater Administration
580 Taylor Avenue
Annapolis, MD 21401
(301) 269-3786

Herbert Austin
VA Institute of Marine
Science
Gloucester Point, VA 23062
(804) 642-7000

Mark Chittenden, Jr.
VA Institute of Marine
Science
Gloucester Point, VA 23062
(804) 642-7000

Dennis Spitsbergen
NC Div. Mar. Fish.
P.O. Box 769
Morehead City, NC 28557
(919) 726-7021

Douglas DeVries
NC Div. Mar. Fish.
P.O. Box 769
Morehead City, NC 28557
(919) 726-7021

Glenn Ulrich
SC Wildl. & Mar. Res. Dept.
P.O. Box 12559
Charleston, SC 29412
(803) 795-6350

James Music
GA Dept. Nat. Resources
Coastal Resources Div.
1200 Glynn Avenue
Brunswick, GA 31523
(912) 264-7221

Roy Williams
FL Dept. Nat. Resources
100 8th Ave., S.E.
St. Petersburg, FL 33701
(813) 896-8626

John Merriner
NMFS-Southeast
Fisheries Center
Beaufort Lab.
Beaufort, NC 28516
(919) 728-4595

Stuart Wilk
NMFS-Northeast
Fisheries Center
Sandy Hook Lab.
Highlands, NJ 07732
(201) 872-0200

Glen Mahoney
NMFS
State Fish Pier
Gloucester, MA 01930
(617) 281-3600

Paul Perra
Atlantic States Marine
Fisheries Commission
1717 Massachusetts Ave.
N. W.
Washington, D.C.
(202) 387-5330

Sciaenid Board

John M. Cronan, Chief
 RI Div. of Fish & Wildlife
 Wash. County Govt. Center
 Tower Hill Road
 Wakefield, RI 02879
 (401) 789-3094

Eric M. Smith
 Assistant Director
 CT Marine Fisheries Office
 PO Box 248
 Waterford, CT 06385
 (203) 442-0166

Gordon Colvin, Director
 NY Div. of Marine & Coastal
 Dept. of Environ. Conservation
 Bldg. 40, SUNY
 Stony Brook, NY 11794
 (516) 751-7900
 Chairman

Bruce Freeman
 Marine Fish. Administrator
 NJ Div. Fish, Game
 and Wildlife
 CN 400
 Trenton, NJ 08625
 (609) 292-2083

L. E. Zeni, Administrator
 MD Tidewater Adm.
 Tawes State Office Building
 Annapolis, MD 21401
 (301) 269-3733

William C. Wagner II, Director
 DE Div. of Fish Wildlife
 Dept. of Natural Res. & Env. Control
 Edward Tatnall Bldg.
 Dover, DE 19901
 (302) 736-5295

William A. Pruitt
 VA Marine Resources Commission
 Box 756
 Newport News, VA 23607

Robert K. Mahood, Director
 NC Div. of Marine Fisheries
 and Community Development
 PO Box 769
 Morehead City, NC 28557
 (919) 726-7021

Edwin B. Joseph, Director
 SC Div. of Marine Resources
 Dept.
 PO Box 12559
 Charleston, SC 29412
 (803) 795-6350

Duane Harris, Director
 GA Coastal Resources Division
 Dept. Nat. Resources
 1200 Glynn Avenue
 Brunswick, GA 31520
 (912) 264-7218

James T. Barrett, Jr.
 Sports Fishing Adm.
 Bur. of Marine Resources
 FL Dept. of Nat. Res.
 3900 Commonwealth Blvd.
 Tallahassee, FL 32303
 (904) 488-1554

John Cookson
 NMFS
 State Fish Pier
 Gloucester, MA
 (617) 281-2600

Irwin Alperin,
 Executive Director
 Atlantic States Marine Fisheries
 Commission
 1717 Massachusetts, NW
 Washington, D.C. 20036
 (202) 387-5330

South Atlantic State/Federal Fisheries Management Board

Robert K. Mahood, Director
NC Div. of Marine Fisheries
Dept. of Natural Resources and
Community Development
PO Box 769
Morehead City, NC 28557
(919) 726-7021

Edwin B. Joseph, Director
SC Div. of Marine Resource
Wildlife & Marine Resources Dept.
PO Box 12559
Charleston, SC 29412
(803) 795-6350

Duane Harris, Director
GA Coastal Resources Div.
Dept. Natural Resources
1200 Glynn Avenue
Brunswick, GA 31520
(912) 264-7218

James T. Barrett, Jr.
Sports Fishing Administrator
Bur. of Marine Resources
FL Dept. of Natural Resources
3900 Commonwealth Blvd.
Tallahassee, FL 32303
(904) 488-1554

I. B. Byrd
National Marine Fisheries Service
Southeast Region
9450 Koger Blvd.
St. Petersburg, FL 33702
(813) 893-3161

Irwin Alperin
Executive Director
Atlantic States Marine
Fisheries Commission
1717 Massachusetts Ave., N.W.
Washington, D.C. 20036
(202) 387-5330

NOAA COASTAL SERVICES CENTER LIBRARY



3 6668 00003 7400